



HAL
open science

Exposure of unweaned calves to the toxic compound of giant fennel

Gilbert Gault, Isabelle Fourel, Sébastien Lefebvre, Virginie Lattard, Denis Grancher

► **To cite this version:**

Gilbert Gault, Isabelle Fourel, Sébastien Lefebvre, Virginie Lattard, Denis Grancher. Exposure of unweaned calves to the toxic compound of giant fennel. Congress of the European Society of Veterinary and Comparative Nutrition, Sep 2017, Cirencester, United Kingdom. , 1 p., 2017. hal-02789023

HAL Id: hal-02789023

<https://hal.inrae.fr/hal-02789023>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Exposure of unweaned calves to the toxic compound of giant fennel

Gilbert Gault, Isabelle Fourel, Sebastien Lefebvre, Virginie Lattard, Denis
Grancher

► **To cite this version:**

Gilbert Gault, Isabelle Fourel, Sebastien Lefebvre, Virginie Lattard, Denis Grancher. Exposure of unweaned calves to the toxic compound of giant fennel. Congress of the European Society of Veterinary and Comparative Nutrition (ESVCN), 2017, Sep 2017, Cirencester, United Kingdom. Congress of the European Society of Veterinary and Comparative Nutrition (ESVCN), Cirencester (England), 2017, 2017. <hal-01590514>

HAL Id: hal-01590514

<https://hal.archives-ouvertes.fr/hal-01590514>

Submitted on 19 Sep 2017

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Gilbert Gault, Isabelle Fourel, Sébastien Lefebvre, Virginie Lattard, Denis Grancher*
 USC 1233 INRA-Vetagro Sup, Veterinary School of Lyon, 1 avenue Bourgelat, 69280 Marcy
 l'Etoile, France
 *denis.grancher@vetagro-sup.fr

Introduction:

- Coagulopathy occurs in animal intoxicated with Giant fennel (*Ferula communis*)
- Giant fennel contains ferulenol which can block vitamin K regeneration cycle by the same mechanism that dicoumarol, which is produced from coumarin in fermented sweet clover hay [1].
- Focus on a Corsica herd of free range cattle exposed to giant fennel
- Does exposition of mother lead to an exposition of unweaned calves ?

Results:

Results are presented in figure 1.

Adults and weaned calves:

30/31 samples showed positive results for ferulenol



Confirm the exposure of cattle to giant fennel

Unweaned calves:

Almost all are positive for ferulenol (4/5)



Source of exposure is milk

Incidental finding :

-The majority of blood samples of adults and weaned calves (28/31) are positive for dicoumarol. The origin of dicoumarol is currently unknown.

Material and methods:





Figure 1: Blood concentration in ferulenol and dicoumarol

Discussion:

- . This study shows that **ferulenol is excreted in milk**. These results raise many questions on the **consequences of these excretions**. Currently, the **absence of reported intoxication in unweaned calves** is consistent with the weak level of ferulenol in their blood (median: 1.1 ng/mL). Further investigations are necessary to assess the possible importance of this exposure in animals and the impact of contaminated milk in **human health**.
- . Regarding dicoumarol, it is not possible to conclude on milk excretion. Moreover, its origin remains a mystery. Sweet clover was not found in hay given by the farmer. The possible presence of dicoumarol in fresh sweet clover growing in the area is under investigation.

Conclusion:

Cows exposed to ferulenol after eating giant fennel excrete ferulenol in milk, that leads to an exposure of unweaned calves.

References:

1. Louvet M-S et al. Comparative inhibitory effect of prenylated coumarins, ferulenol and ferprenin, contained in the 'poisonous chemotype' of *Ferula communis* on mammal liver microsomal VKORC1 activity. *Phytochemistry*. oct 2015;118:124-30.
2. Fourel I et al. Validation of a new liquid chromatography- tandem mass spectrometry ion-trap technique for the simultaneous determination of thirteen anticoagulant rodenticides, drugs, or natural products. *J Anal Toxicol*. mars 2010;34(2):95-102.

All publications of USC 1233 are freely available on HAL

