

# Survey of genetic diversity in European Populus breeding programs

Véronique Jorge, Arnaud Dowkiw, Patricia Faivre-Rampant, Marc Villar, Marie Pegard, Vincent Segura, Vanina Guérin, Marijke Steenackers, Lorenzo Vietto, Catherine Bastien

#### ▶ To cite this version:

Véronique Jorge, Arnaud Dowkiw, Patricia Faivre-Rampant, Marc Villar, Marie Pegard, et al.. Survey of genetic diversity in European Populus breeding programs. INUPRAG Meeting 2018, Jun 2018, Barcelone, Spain. hal-02789044

#### HAL Id: hal-02789044 https://hal.inrae.fr/hal-02789044v1

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





# Survey of genetic diversity in European Populus breeding programs

**CONTRIBUTORS** to the present work:

V. Jorge, A. Dowkiw, P. Faivre-Rampant, M. Villar, M. Pegard, V. Segura, V. Guérin, M.

Steenackers, L. Vietto, C. Bastien INRA Experimental unit GBFOR








#### Our research objective

Management of forest genetic resources for sustainable timber production in a changing climate context

Individuals

**Populations** 

Management Units (Conservation, Breeding)



Genetic Resources for research

in- & ex-situ Conservation

Breeding programmes

**Operational Pole** « Genetic Resources »

> FRM production engineering.


Guidelines for FRM Use



Joint unit between **INRA and ONF** (French National Forest Service)



- ~50 persons (40 perm.)
- 4 teams



**Species studied :** Poplars, Douglas Fir, Scot Pine, wild cherry, larches, ash





# SUMMARY

- - 1. Poplars and poplar cultivars
  - 2. Objectives of the study
  - 3. Material and methods
  - 4. Genotyping quality and discrimination between species
  - 5. Diversity of genitors used in poplar breeding programs
  - 6. Relatedness between cultivars
  - 7. Diversity in multi-clonal varieties
  - 8. Conclusions



# Poplars (genus Populus)

#### 6 sections et 29 species Dioecious



Populus nigra var 'Italica'









Populus trichocarpa Torr. & A. Gray




Populus deltoides Bartr. ex Marsh.




# **Breeding poplars**

- > Recurrent selection at intraspecific level
- > Interspecific hybrids => hybrid vigour





# Poplar cultivation and consequences



Figures from TreeBreedeX Monography contributions

#### List of registered material (2014)

- > 172 monoclonal cultivars
- > 10 multi-clonal varieties

But only few cultivars used by European countries ...


# Nb of clones representing 90% of planting stock (2007):

France: 9 clones, I-214 (24%) - Triplo (10%) Italy: 7 clones, I-214 (77%) - Villafranca (3%) Belgium: 16 clones, Koster (23%) - Unal (1%)

Spain: 5 clones, I-214 (70%)



- Monoclonal stands
- Few cultivars used with « fashion » effect
- Diseases outbreaks in a P. deltoides x P. trichocarpa context





©S. Bisoff








- Cultivated (exotic, ornemental clones) and native poplars often coexist in the landscape
- > Introgression as a potential risk for natural resources



- Cultivated (exotic, ornemental clones) and native poplars often coexist in the landscape
- > Introgression as a potential risk for natural resources



Black poplar






- > Cultivated (exotic, ornemental clones) and native poplars often coexist in the landscape
- Introgression as a potential risk for natural resources



Black poplar





Lombardy poplar « Italica »






- Cultivated (exotic, ornemental clones) and native poplars often coexist in the landscape
- > Introgression as a potential risk for natural resources



Black poplar





Lombardy poplar
« Italica »



cultivated poplars stands





# 2. Objectives of the study



## Gentree (H2020) project – WP2

Optimising the management and sustainable use of forest genetic resources in Europe

WP2 – Innovative breeding strategies based on the rational use of forest genetic resources

#### Task 2.2:



Design and Implementation of allele and genotype sampling strategies to enrich breeding populations

- Comparative analysis of genetic diversity in natural / breeding populations: Populus nigra in EU breeding programmes
- ➤ P. deltoides x P. nigra cultivars with high tolerance to pests and diseases and high productivity even in northern conditions: take value of large genetic diversity available from natural P. nigra ressources





# 3. Material and methods



#### **Plant material**

- Cultivars incl. 'Bellini', Italian cultivar!
- Genitors from active breeding programs (BE, FR, IT)
- Multi-Clonal Varieties











| type                                   | origin         | #                          |
|--|----------------|----------------------------|
| Cultivars <i>P. nigra</i> , ornemental |                | Italica, Thevestina        |
| Cultivars P. nigra registered          | F/I, NL        | Jean Pourtet,<br>Vereecken |
| Cultivars P. x<br>canadensis           | Belgium        | 5                          |
|  | France         | 5 (old cultivars)          |
|  | Hungary        | 2                          |
|  | Italy          | 29                         |
|  | NL             | 5                          |
|  | Poloni         | 6                          |
| Genitors                               | Alasia (Italy) | 10 (DN)                    |
|  | CREA (Italy)   | 28                         |
|  | GIS (France)   | 39                         |
|  | INBO (Belgium) | 11 (6 DN; 5 N)             |
| Multiclonal<br>Varieties (MCV)         | France         | 2 (25 ind. each)           |



# 10 K SNP array used characterise genetic diversity in *P. nigra* at European scale



- Diversity structured according to drainage systems.
- An ancestral cluster clearly admixed in <u>almost all</u> predefined populations.
- Introgression from cultivated stands?






# 4. Genotyping quality and discrimination between species




4. Genotyping quality and discrimination between species

# **Genotyping quality**



- Quality of initial genotyping: 11% SNP failed => 9 127 SNPs to use
- Overlap with 2 previous genotyping campaigns including natural populations => 6 298 SNPs
- Selection of SNP to answer different questions :
  - > ~1 200 SNP selected evenly distributed on the genome to describe diversity in poplar breeding programs.
  - Remark: Only 802 SNP showed 3 genotypic classes in the set of P. deltoides x P. nigra cultivars => those SNP polymorphic in P. deltoides.

# Discrimination between species



- Potential of the genotyping tool to discriminate *Populus* species and hybrids (axe 1).
- Impact of ornamental *Populus nigra* cv (Italica, axe 2).

#### **Perspectives:**

Development of discriminant markers for axe 1 and axe 2

« hybrid » cultivars + others DN, DNxN P. nigra genitors

+ VMC






# 5. Diversity of genitors used in poplar breeding programs



# Large diversity and low introgression / Italica, except for Belgium breeding population

P. nigra genitors vs. Natural populations (in grey)
PCA axes 1 & 2, 1128 SNP






# 6. Relatedness between cultivars



#### 5. Relatedness between cultivars

#### **Genetic distances**



- Confirmation of previous conclusions: distinction between taxa, genitors pooled/breeding programs
- > Different levels of relatedness inside breeding programs
- Close relationship between some cultivars
- Pure P. deltoides among declared P. deltoides x P. nigra hybrids ?

1128 SNPs

Belgium

France (GIS, Poloni)

Italy


Netherland

+ old cultivars



# **Evidence of the use of Italica in poplar breeding**

Resolving pedigree of some culitvars.



Results from Colony, Jones & Wang 2010






# 6. Diversity in multiclonal varieties



### Multi-Clonal Varieties of Populus nigra

#### Target in deployment = ecological restoration of river banks



#### General rules of a MCV of black poplar

- Individuals of the species Populus nigra!
- Variety by river basin (six in France)
- Largest diversity as possible (geographic, ecologic and genetic), representative of the river basin
- Choice of 25 genotypes per variety

Occurence of the species:

Present in riparian forests

Present as isolated trees

absent

Development of cost- and time-effective breeding methods = T2.3 of GenTree project



### Multi-Clonal Varieties of Populus nigra

#### **Actual methodology**

#### Species descriptors and introgression

- In situ phenotype : <u>tree shape</u>, bark characteristics, stem characteristics, leaf shape, bud shape, male and female catkins
- SSR markers

# One variety per river basin & large ecological regions

GRECO: 'large ecological regions' of the French National Forest Inventory



Populus nigra var. Italica



#### Largest diversity as possible

- Sex ratio
- No clone (SSR markers)

Traits evaluated in trials:

- Phenology: max. variability
- Height and diameter: eliminate extremes to avoid competiton
- Branching patterns: avoid fastigiate







### Multi-Clonal Varieties of Populus nigra

- Registration on French national register of 2 new multi-clonal varieties of *Populus nigra* for the Rhône river basin
- Validation of the low-cost strategy with SNP genotyping




Populus nigra multiclonal variety: 2 yrs-old plants

1128 SNPs

**Grey**: Rhone natural populations

**Greens: individuals selected for the MCV** 



# 8. Conclusions



#### **Conclusions**

- ➤ 10K SNP array valuable tool to evaluate diversity in *P. nigra* but limited for *P. deltoides* (but not nul?)
- ➤ Actual breeding programs used local *P. nigra* diversity => but hybrid cultivars bred are registered at European level
- Validation of a low-cost selection methodology for MCV



How to design and implement allele and genotype sampling strategies to enrich breeding populations?

#### GenTree Task 2.4

Improvement of genetic diversity management in intensive breeding programmes

Genomic Selection
explicite management of genetic diversity
Optimisation introduction new diversity





# Graciès!

#### **Projects**







**INRA Selgen project** Breed2Last



#### **Partners**











INSTITUUT NATUUR- EN BOSONDERZOEK



