



## Survey of genetic diversity in European Populus breeding programs

Véronique Jorge, Arnaud Dowkiw, Patricia Faivre-Rampant, Marc Villar, Marie Pegard, Vincent Segura, Vanina Guérin, Marijke Steenackers, Lorenzo Vietto, Catherine Bastien

### ► To cite this version:

Véronique Jorge, Arnaud Dowkiw, Patricia Faivre-Rampant, Marc Villar, Marie Pegard, et al.. Survey of genetic diversity in European Populus breeding programs. INUPRAG Meeting 2018, Jun 2018, Barcelone, Spain. hal-02789044

**HAL Id: hal-02789044**

**<https://hal.inrae.fr/hal-02789044>**

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



BioForA



©S. Bisoffi

# Survey of genetic diversity in European Populus breeding programs


CONTRIBUTORS to the present work :

V. Jorge, A. Dowkiw, P. Faivre-Rampant, M. Villar, M. Pegard, V. Segura, V. Guérin, M. Steenackers, L. Vietto, C. Bastien  
INRA Experimental unit GBFOR



## Our research objective


Management of forest genetic resources for sustainable timber production in a changing climate context



- **Joint unit between INRA and ONF** (French National Forest Service)




- **~50 persons (40 perm.)**
- **4 teams**



**Species studied :** Poplars, Douglas Fir, Scot Pine, wild cherry, larches, ash

# SUMMARY

1. **Poplars and poplar cultivars**
2. **Objectives of the study**
3. **Material and methods**
4. **Genotyping quality and discrimination between species**
5. **Diversity of genitors used in poplar breeding programs**
6. **Relatedness between cultivars**
7. **Diversity in multi-clonal varieties**
8. **Conclusions**



# 1. Poplars and poplar cultivars



## Poplars (genus *Populus*)

6 sections et 29 species

Dioecious



*Populus nigra* var 'Italica'



*Populus nigra* L.



*Populus alba* L.



*Populus tremula* L.




*Populus trichocarpa* Torr. & A. Gray




*Populus deltoides* Bartr. ex Marsh.

## Breeding poplars

- Recurrent selection at intraspecific level
- Interspecific hybrids => hybrid vigour



## Poplar cultivation and consequences



Figures from TreeBreedX Monography contributions

### List of registered material (2014)

- 172 monoclonal cultivars
- 10 multi-clonal varieties

*But only few cultivars used by European countries ...*

### Nb of clones representing 90% of planting stock (2007) :


France : 9 clones, I-214 (24%) - Triplo (10%)  
Italy : 7 clones, I-214 (77%) – Villafranca (3%)  
Belgium : 16 clones, Koster (23%) - Unal (1%)  
Spain : 5 clones, I-214 (70%)





## Poplar cultivation and consequences

- Monoclonal stands
- Few cultivars used with « fashion » effect
- Diseases outbreaks in a *P. deltoides* x *P. trichocarpa* context



©S. Bisoffi



# Poplar cultivation and consequences

- Cultivated (exotic, ornamental clones) and native poplars often coexist in the landscape
- Introgression as a potential risk for natural resources



© C. Maître, INRA




# Poplar cultivation and consequences

- Cultivated (exotic, ornamental clones) and native poplars often coexist in the landscape
- Introgression as a potential risk for natural resources



***Black poplar***




© C. Maître, INRA

## Poplar cultivation and consequences

- Cultivated (exotic, ornamental clones) and native poplars often coexist in the landscape
- Introgression as a potential risk for natural resources



***Black poplar***



***Lombardy poplar***  
« Italica »

© C. Maître, INRA




## Poplar cultivation and consequences

- Cultivated (exotic, ornamental clones) and native poplars often coexist in the landscape
- Introgression as a potential risk for natural resources



**Black poplar**




**Lombardy poplar**  
« Italica »



**cultivated  
poplars  
stands**

© C. Maître, INRA



## 2. Objectives of the study

# Gentree (H2020) project – WP2

Optimising the management and sustainable use of forest genetic resources in Europe


**WP2** – Innovative breeding strategies based on the rational use of forest genetic resources


## Task 2.2:



Design and Implementation of allele and genotype sampling strategies to enrich breeding populations

- Comparative analysis of genetic diversity in natural / breeding populations: *Populus nigra* in EU breeding programmes
- *P. deltoides* x *P. nigra* cultivars with high tolerance to pests and diseases and high productivity even in northern conditions : take value of large genetic diversity available from natural *P. nigra* resources





# 3. Material and methods



## Plant material

#### ➤ Cultivars

incl. 'Bellini', Italian cultivar !


#### ➤ Genitors from active breeding programs (BE, FR, IT)

#### ➤ Multi-Clonal Varieties





type	origin	#
Cultivars <i>P. nigra</i> , ornamental		Italica, Thevestina
Cultivars <i>P. nigra</i> registered	F/I, NL	Jean Pourtet, Vereecken
Cultivars <i>P. x canadensis</i>	Belgium	5
	France	5 (old cultivars)
	Hungary	2
	Italy	29
	NL	5
	Poloni	6
Genitors	Alasia (Italy)	10 (DN)
	CREA (Italy)	28
	GIS (France)	39
	INBO (Belgium)	11 (6 DN; 5 N)
Multiclonal Varieties (MCV)	France	2 (25 ind. each)

## 10 K SNP array used to characterise genetic diversity in *P. nigra* at European scale




- ❖ Diversity structured according to drainage systems.
- ❖ An **ancestral** cluster clearly admixed in almost all predefined populations.
- ❖ Introgression from cultivated stands ?





## 4. Genotyping quality and discrimination between species


## Genotyping quality



- Quality of initial genotyping : **11% SNP failed** => 9 127 SNPs to use
- Overlap with 2 previous genotyping campaigns including natural populations => **6 298 SNPs**
- Selection of SNP to answer different questions :
  - **~1 200 SNP** selected evenly distributed on the genome to describe diversity in poplar breeding programs.
  - Remark : Only **802 SNP** showed 3 genotypic classes in the set of *P. deltoides* x *P. nigra* cultivars => those SNP polymorphic in *P. deltoides*.



# Discrimination between species




- Potential of the genotyping tool to discriminate *Populus* species and hybrids (axe 1).
- Impact of ornamental *Populus nigra* cv (Italica, axe 2).

## Perspectives :

Development of discriminant markers for axe 1 and axe 2

« hybrid » cultivars  
+ others DN, DNxN


*P. nigra* genitors  
+ VMC




## 5. Diversity of genitors used in poplar breeding programs

# Large diversity and low introgression / Italica, except for Belgium breeding population

*P. nigra* genitors vs. Natural populations (in grey)  
PCA axes 1 & 2, 1128 SNP




- Each breeding population representative of the local natural population
- How much this diversity is representative ?
- Perspectives: compare effective population size.



## 6. Relatedness between cultivars



# Genetic distances




- Confirmation of previous conclusions : distinction between taxa, genitors pooled/breeding programs
- Different levels of relatedness inside breeding programs
- Close relationship between some cultivars
- Pure *P. deltooides* among declared *P. deltooides* x *P. nigra* hybrids ?


1128 SNPs  
Belgium  
France (GIS, Poloni)  
Italy  
Netherland  
+ old cultivars

# Evidence of the use of *Italica* in poplar breeding

❖ Resolving pedigree of some cultivars.







*Results from Colony, Jones & Wang 2010*



## 6. Diversity in multiclonal varieties

## Target in deployment



 Present in riparian forests  
 Present as isolated trees  
 absent

- Individuals of the species *Populus nigra* !
- Variety by river basin (six in France)
- Largest diversity as possible (geographic, ecologic and genetic), representative of the river basin
- Choice of 25 genotypes per variety



## Multi-Clonal Varieties of *Populus nigra*


### Actual methodology

#### Species descriptors and introgression

- In situ phenotype : tree shape, bark characteristics, stem characteristics, leaf shape, bud shape, male and female catkins
- SSR markers

#### One variety per river basin & large ecological regions

GRECO : 'large ecological regions' of the French National Forest Inventory



*Populus nigra* var. *Italica*



#### Largest diversity as possible

- Sex ratio
  - No clone (SSR markers)
- Traits evaluated in trials :
- Phenology: max. variability
  - Height and diameter: eliminate extremes to avoid competition
  - Branching patterns: avoid fastigate



## Multi-Clonal Varieties of *Populus nigra*

- **Registration** on French national register of 2 new multi-clonal varieties of *Populus nigra* for the Rhône river basin
- Validation of the low-cost strategy with SNP genotyping





*Populus nigra* multiclonal variety: 2 yrs-old plants

1128 SNPs

Grey : Rhone natural populations

Greens : individuals selected for the MCV





## 8. Conclusions

# Conclusions

- 10K SNP array valuable tool to evaluate diversity in *P. nigra* but limited for *P. deltoides* (but not nul ?)
- Actual breeding programs used local *P. nigra* diversity => but hybrid cultivars bred are registered at European level
- Validation of a low-cost selection methodology for MCV



How to design and implement allele and genotype sampling strategies to enrich breeding populations ?

## GenTree Task 2.4

Improvement of genetic diversity  
management in intensive breeding  
programmes

Genomic Selection  
explicite management of genetic diversity  
Optimisation introduction new diversity



# Graciès !

## Projects



INRA Selgen project  
Breed2Last



## Partners

GIS peuplier



INSTITUUT  
NATUUR- EN  
BOSONDERZOEK



Vlaanderen  
is wetenschap