



HAL
open science

GWAS of digestive efficiency in chicken

Sandrine Mignon-Grasteau, Christelle Hennequet-Antier, Elisabeth Le Bihan-Duval

► **To cite this version:**

Sandrine Mignon-Grasteau, Christelle Hennequet-Antier, Elisabeth Le Bihan-Duval. GWAS of digestive efficiency in chicken. 3. Annual Meeting of H2020 Feed-a-Gene project, Apr 2018, Newcastle, United Kingdom. 1 p., 2018. <hal-02789047>

HAL Id: hal-02789047

<https://hal.inrae.fr/hal-02789047v1>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization



GWAS of digestive efficiency in chicken

Sandrine Mignon-Grasteau¹, Christelle Hennequet-Antier¹,
Elisabeth Le Bihan-Duval¹

1. BOA, Institut national de la recherche agronomique (INRA), Université de Tours, 37380 Nouzilly, France.

Objectives

Feedstuffs used in poultry diets are more and more varied to cope with the increased cost of usual high quality feedstuffs and to limit environmental impact of animal productions. As these new feedstuffs are not as optimal as corn or soybean, animals are more challenged and have to be able to adapt to these new feeding conditions. Digestive efficiency is an essential component of chicken adaptability to feed. Heritability estimates of digestive efficiency indicated that this trait is partly under genetic control (Mignon-Grasteau et al., 2004). The purpose of this work was to improve our knowledge on the genetic basis of digestive efficiency through a genome-wide association study on chickens selected for their digestive efficiency.

Materials and methods

Animals



We used 192 animals from the 8th generation of an advanced intercross line between two divergent lines of broilers selected for their high or low digestive efficiency (criterion of selection: AMEn at 3 weeks, using a wheat-based diet). Before the start of cross, the difference in AMEn between the two divergent lines was around 30-40%.

Birds were reared on floor from hatch to 11 days and then transferred to individual cages for digestive efficiency measurement. Animals were fed with a wheat-based diet containing 55% of Rialto wheat, a viscous and hard variety.

Balance trial

At 3 weeks, a balance trial was performed, using the method of total collection of feces of Bourdillon et al. (1990). After freeze-drying, feces were analyzed by NIRS to obtain AMEn, and coefficients of digestive use of dry matter, starch and nitrogen and lipids (Bastianelli et al., 2010).

Genetic analyses

At the end of balance trial, animals were blood sampled and genotyped with the 540K SNP Affymetrix® Axiom® chip. After quality control (MAF, Hardy-Weinberg, Call rate), 353888 SNP were retained for the analysis. These SNP were located on chromosomes 1 to 28 and 33. Phenotypic data were pre-corrected for significant fixed effects. GEMMA software was used for GWAS analyses, with an univariate linear mixed model (polygenic effect, SNP effect). A Bonferroni correction was applied at genome level to take into account multiple testing.

GWAS results

A total of 12 SNPs were genome-wide significant (Bonferroni adjusted *P*-value < 0.05) for AMEn, coefficients of digestive use of dry matter, starch and nitrogen. None was significant for coefficient of digestive use of lipids.

Figure 1. Manhattan plot for AMEn




Figure 2. Manhattan plot for coefficient of digestive use of starch




Table 1. Genes associated with 5% genome-wide significant SNPs.

Chr	SNP	Trait	Within Gene/ Intergenic
1	144 284 848	CDU (DM, Starch, Nitrogen)	ZIC2
2	41 039 328	CDU (DM, Starch, Nitrogen)	ANO10
2	116 692 708	AMEn, CDU (DM, Starch, Nitrogen)	SULF1
3	81 690 202	AMEn, CDU (DM, Starch, Nitrogen)	CD109
4	85 583 360	CDU (DM, Starch, Nitrogen)	Intergenic
5	23 444 968	AMEn, CDU (DM, Starch, Nitrogen)	F2
8	26 671 543	CDU (Starch, Nitrogen)	FGGY
19	6 390 258	CDU (Starch, Nitrogen)	Intergenic
19	6 677 286	CDU (Starch, Nitrogen)	Intergenic
19	6 764 624	CDU (Starch, Nitrogen)	Intergenic
21	897 100	AMEn, CDU (Starch, Nitrogen)	1000 bp before C1ORF174
26	2 317 884	CDU (Starch, Nitrogen)	PM20D1

Conclusions

These results suggest a list of potential candidate genes involved in digestive efficiency. This list will be enlarged by GWAS analyses on related phenotypes measured on the same birds (feed intake, feed conversion ratio, anatomy of digestive tract) and with transcriptomic data (gizzard, jejunum) obtained on a sample of birds from the F2 generation.

References

- Bastianelli D, Bonnal L, Juin H, Mignon-Grasteau S, Davrieux F, Carré B. J NIRS 18:69-77.
- Bourdillon A, Carré B, Conan L, Duperray J, Huyghebaert G, Leclercq B, Lessire M, McNab J, Wiseman J. 1990. Br Poult Sci. 31:557-65.
- Mignon-Grasteau S, Muley N., Bastianelli D, Gomez J, Péron A, Sellier N, Millet N, Besnard J, Hallouis JM, Carré B. 2004. Poult Sci. 83 : 860-867.



www.feed-a-gene.eu



The Feed-a-Gene Project has received funding from the European Union's H2020 Programme under grant agreement no 633531.