



The pace of aggressiveness evolution and host adaptation in a plant pathogen population

Frederic Suffert, Henriette Goyeau, Ivan Sache, Florence Carpentier, Sandrine Gélisse, David Morais

► To cite this version:

Frederic Suffert, Henriette Goyeau, Ivan Sache, Florence Carpentier, Sandrine Gélisse, et al.. The pace of aggressiveness evolution and host adaptation in a plant pathogen population. Journées Jean Chevau-
chon JJC2018 - 12èmes Rencontres de Phytopathologie & Mycologie,, Jan 2018, Aussois, France. p.34.
hal-02791218

HAL Id: hal-02791218

<https://hal.inrae.fr/hal-02791218>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

EP08

The pace of aggressiveness evolution and host adaptation in a plant pathogen population

Frédéric Suffert, Henriette Goyeau, Ivan Sâche, Florence Carpentier, Sandrine Gélisse, David Morais, Ghislain Delestre

UMR BIOGER, INRA, AgroParisTech, Université Paris-Saclay, 78850 Thiverval-Grignon, France.

The efficiency of plant resistance to fungal pathogen populations is expected to decrease over time, due to an increase in the frequency of virulent, or highly aggressive strains. This dynamics may differ depending on time scale (annual or pluriannual), especially for annual crop pathogens with both sexual and asexual reproduction modes. In this study we investigated the pace of pathogen evolution and host adaptation. To this end we compared changes in two aggressiveness traits, latent period and lesion size, in a *Zymoseptoria tritici* population, over an eight-month growing season and a six-year period of wheat monoculture. Two pairs of subpopulations were collected in a wheat plot, the first at the beginning and end of a single annual epidemic, and the second at the beginning and end of the six-year period. We established a trend toward decreased latent period during the course of the annual epidemic, concomitant with a significant loss of variability, but not over the six-year period. We interpreted this contrast as the consequence of a trade-off between, on the one hand, a gain in aggressiveness by selection after several generations of asexual reproduction at the annual scale and, on the other hand, a decrease in the probability of sexual reproduction (Allee effect) and transmission to the next epidemic season. The conclusions of this study should be taken into account in the design and test of scenarios of deployment of partial resistance at the landscape scale.