



## Exploring the genetic variability of peach skin phenolics and triterpenoids as natural defenses against brown rot

Bénédicte Quilot-Turion, Silvia Scariotto, Marie Noëlle Corre, Leandro de Oliveira-Lino, Véronique Signoret, Maximiliano Dini Vinoly, Lorraine Latchoumane, Maria Do Carmo Raseira, Jean-Luc Poëssel

### ► To cite this version:

Bénédicte Quilot-Turion, Silvia Scariotto, Marie Noëlle Corre, Leandro de Oliveira-Lino, Véronique Signoret, et al.. Exploring the genetic variability of peach skin phenolics and triterpenoids as natural defenses against brown rot. 9. International Rosaceae Genomics Conference, Nanjing Agricultural University (NAU). Nanjing, CHN., Jun 2018, Nanjing, China. hal-02791339

**HAL Id: hal-02791339**

**<https://hal.inrae.fr/hal-02791339>**

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Exploring the genetic variability of peach skin phenolics and triterpenoids as natural defenses against brown rot

QUILOT-TURION Bénédicte<sup>1§</sup>, SCARIOTTO Silvia<sup>1,2</sup>, CORRE Marie-Noëlle<sup>1</sup>, OLIVEIRA LINO Leandro<sup>1</sup>, SIGNORET Véronique<sup>1</sup>, DINI VINOLY Maximiliano<sup>2</sup>, LATCHOUMANE Lorraine<sup>1</sup>, RASEIRA Maria Do Carmo<sup>2</sup>, POËSSEL Jean-Luc<sup>1</sup>  
(<sup>1</sup>GAFL, INRA, 84000, Avignon, France ; <sup>2</sup>EMBRAPA Clima Temperado, Pelotas, RS, 96010-971, BRAZIL)

<sup>§</sup>benedicte.quilot-turion@inra.fr

## Abstract:

Brown rot caused by *Monilinia* spp. provokes dramatic losses of peach in all production regions. In this context, the general objective of our work is to develop sustainable ways to fight against this damaging disease.

This study explored phenolic and triterpenoid diversity of fruit skin together with brown rot resistance. A Brazilian peach collection and an interspecific cross between *P. davidiana* and *P. persica*, each of 120 genotyped individuals, have been screened for 3 years, for susceptibility to *M. fructicola* and to *M. laxa* respectively. Some genotypes of the two populations showed low or null infection the 3 years.

Secondary compounds of fruit skin of all genotypes were analyzed two years by HPLC-DAD. Depending on the genotypes, 30 to 40 compounds including triterpenoids, hydroxycinnamic derivatives and flavonoids, were quantified and around 20 were identified by mass spectrometry. Among these, phenolic esters of triterpenoids were identified for the first time in peach fruit.

The correlation of quantitative data between years showed their robustness. A large diversity in contents of the compounds was observed.

The association and QTL analyses proved to be complementary and led to assumptions about the co-locations of loci controlling different traits, especially for compounds with differential levels in the skin of peaches and nectarines. These analyses will help deciphering the genetic control of fruit skin phenolics and triterpenoids and pave the way to the identification of the underlying genes. Detection of loci related to resistance will provide tools for the implementation of marker assisted selection.

Correlations between compounds and infection traits led us to hypothesize that some compounds from peach skin could play a role in the brown rot control. The fungicide activity of these compounds is being tested *in vitro*.

*This project is supported by Agropolis Fondation reference ID 1503-003 \_ ANR-10-LABX-0001-01*

**Key words:** Disease resistance, QTL, GWAS, *Monilinia* spp., *Prunus persica*