



**HAL**  
open science

## What do models tell us about water and sediment connectivity ?

Jantiene Baartman, Rens Masselink, João Pedro Nunes, Frédéric Darboux, Charles Bielders, Vincent Cantreuil, Aurore Degré, Petr Kavka, Olivier Cerdan, Thomas Grangeon, et al.

### ► To cite this version:

Jantiene Baartman, Rens Masselink, João Pedro Nunes, Frédéric Darboux, Charles Bielders, et al.. What do models tell us about water and sediment connectivity ?. EGU, European Geosciences Union General Assembly 2018, European Geosciences Union (EGU). DEU., Apr 2018, Vienne, Austria. hal-02791594

**HAL Id: hal-02791594**

**<https://hal.inrae.fr/hal-02791594>**

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Distributed under a Creative Commons Attribution - NonCommercial - NoDerivatives 4.0 International License



## What do models tell us about water and sediment connectivity?

Jantiene Baartman (1), Rens Masselink (1), Joao Pedro Nunes (2), Frederic Darboux (3), Charles Biolders (4), Vincent Cantreul (5), Aurore Degré (5), Petr Kavka (6), Olivier Cerdan (7), Thomas Grangeon (7), Peter Fiener (8), Florian Wilken (8), Marcus Schindewolf (9), Laura Turnbull (10), and John Wainwright (10)

(1) Wageningen University, Soil Physics and Land Management Group, Wageningen, Netherlands (jantiene.baartman@wur.nl), (2) CE3C – Centre for Ecology, Evolution and Environmental Changes, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal, (3) INRA, Laboratoire Sols et Environnement, UMR 1120, Vandoeuvre-lès-Nancy, France, (4) Earth and Life Institute - Environmental Sciences and Faculty of Bioscience Engineering, Université catholique de Louvain, B-1348 Louvain-la-Neuve, Belgium, (5) ULiege, Gembloux Agro-Bio Tech, BIOSE, Passage des Déportés 2, 5030 Gembloux, Belgium, (6) Czech Technical University, Faculty of Civil Engineering, Department of irrigation, drainage and landscape engineering, Prague 6, Czech Republic, (7) BRGM, DRP/RIG, Orléans, France, (8) Institute of Geography, University Augsburg, Augsburg, Germany, (9) Institute of Geography, Catholic University of Eichstätt-Ingolstadt, Ostenstrasse 18, 85072 Eichstätt, Germany, (10) Department of Geography, Science Laboratories, Durham University, Durham, UK

Recently, connectivity has emerged as a promising concept to understand the transfer of water and sediment in a catchment. Both structural connectivity – i.e. representing the connectivity of system properties such as the (micro)topography, and functional connectivity – i.e. representing connections that may change and evolve over time such as soil moisture, are important to consider. As discussed by Nunes et al. (in press), good models should be effectively connected models, i.e. represent properly the fluxes of water and sediment both within and between its fundamental spatial units. However, there is no clear framework to guide how this should be assessed. In this study we analysed changes in landscape connectivity using six well-known erosion models: Erosion3D, Fullswof, Landsoil, LISEM, MAHLERAN and Watersed. Our objective was to determine if, and how simulated connectivity is linked with model structure. The models all simulated the same, semi-virtual 124 ha watershed, loosely based on the Giser experimental agricultural watershed in Belgium. A total of 53 connectivity scenarios with differences in spatial complexity and presence of connectivity features were simulated using two rainfall events (10- and 50-year return periods). The spatial complexity was varied in terms of field size (5, 10 or 20 ha) and five different land-use patterns with tillage orientation following the axis of the fields. Finally, for mid-sized fields (10 ha) and for each of the five land-use patterns, the following connectivity features were tested: conservation tillage (i.e. orientation of fields along the contour), and presence of grass strips and a grassed waterway. We will discuss the impacts of these scenarios on overland flow and sediment connectivity for each model, in relation with its structure. In the future, these results will be used to investigate the possibility to derive more generic results using model ensembles.

### Reference:

Nunes JP, Wainwright J, Biolders CL, Darboux F, Fiener P, Finger D, Turnbull L. In Press. Better models are more effectively connected models. *Earth Surface Processes and Landforms* doi: 10.1002/esp.4323.