



HAL
open science

Detecting molecular footprint of local adaptation in *Zymoseptoria tritici*

Anne Genissel, Lilian L. Gout

► **To cite this version:**

Anne Genissel, Lilian L. Gout. Detecting molecular footprint of local adaptation in *Zymoseptoria tritici*. ECFG13 European conference on fungal genetics, Apr 2016, PARIS LA VILLETTE, France. p.509, 2016. <hal-02792403>

HAL Id: hal-02792403

<https://hal.inrae.fr/hal-02792403v1>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

POSTER SESSION ABSTRACTS
Session CS6 Ecological and population genomics
CS6W38

Wednesday 6th April
14:00 - 16:00

GENISSEL Anne (1), GOUT Lilian (2)

(1) BIOGER, INRA, , AgroParisTech, Thiverval-Grignon, France

(2) BIOGER, AgroParisTech, INRA, Thiverval-Grignon, France

Detecting molecular footprint of local adaptation in *Zymoseptoria tritici*

Zymoseptoria tritici (Zt) is one of the most common pathogen of wheat and yet the genetic basis of adaptation in this species is largely unknown. Population genetics studies using neutral markers suggest for the most part that there is a low level of genetic differentiation between natural population samples in Zt. Using new generation sequencing data, now we can provide an in depth analysis of the genetic variation segregating in natural populations, and better understand the mode and tempo of genome evolution at the intraspecific level. The aim of this work was to identify the genes / genome regions that are under extreme genetic differentiation between two French population samples that differ in environment, collected in the North and South of France. To reach this goal, we used new generation sequencing data from 30 isolates. We found a large genetic diversity for both single nucleotide polymorphisms and structural variants within and between population samples. Many polymorphic sites throughout the genome were highly differentiated between the population samples, especially we discovered a large inversion on chromosome 7. We detailed the nature of genetic variants in this specific region. We also described the pattern of linkage disequilibrium and quantify the skew in allele frequency spectrum across inverted regions to understand the nature of the selection in action. Last, we discuss the potential role of inversions in local adaptation.
