



**HAL**  
open science

## **Bacillus cereus: diversity and adaptation to conditions in the food chain**

Frederic F. Carlin

► **To cite this version:**

Frederic F. Carlin. Bacillus cereus: diversity and adaptation to conditions in the food chain. IAFP European Symposium on Food Safety, Apr 2015, Cardiff, United Kingdom. hal-02792444

**HAL Id: hal-02792444**

**<https://hal.inrae.fr/hal-02792444>**

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

*Bacillus cereus*: diversity and adaptation to conditions in the food chain.

Frédéric CARLIN

INRA, Université d'Avignon, UMR408 Sécurité et Qualité des Produits d'Origine Végétale,  
Site Agroparc, 84914 Avignon cedex 9, France

[frederic.carlin@avignon.inra.fr](mailto:frederic.carlin@avignon.inra.fr)

*Bacillus cereus* accounts for some of the most common foodborne poisonings. Reports of *Bacillus cereus* outbreaks in Europe (including very severe emetic outbreaks) markedly increased in recent years. Spores of *B. cereus* are widely dispersed in the environment and contaminate any sort of foods. With *Clostridium botulinum* and *C. perfringens*, *B. cereus* is a concern for mildly heat-processed foods because of spore resistance and many strains have the ability to multiply at refrigeration temperature. The phenotypic diversity of *Bacillus cereus* strains has been established long ago. Recently a robust description of the genetic structure of the whole group has been provided and defined seven genetic groups. With respect to food safety, these groups numbered I-VII have different growth temperature profiles, and also differ by their involvement in foodborne poisoning incidents, production of toxins causing poisonings, resistance to heat, adaptation to low pH or to low  $a_w$ , etc... For instance most cereulide-producing strains are distributed in phylogenetic group III that also contains the most heat-resistant strains. The "diarrhoeic potential" is low for psychrotrophic group VI strains, while it is higher for mesophilic groups III and IV, which is consistent with the prevalence of foodborne disease strains in the groups. A procedure to identify a *B. cereus* strain in the seven phylogenetic groups has been proposed. The risk for consumers of *B. cereus* foodborne poisonings should be considered with respect to the type of strains present in foods.