



HAL
open science

Survey of fungal lignocellulose-acting enzymes for improved plant biomass degradation capacities

Eric Record, Anne Lomascolo, Anne Favel, David D. Navarro, Laurence L. Lesage-Meessen, Annick Doan, Mireille M. Haon, Christophe Boyer, Sabine Taussac, Sacha S. Grisel, et al.

► To cite this version:

Eric Record, Anne Lomascolo, Anne Favel, David D. Navarro, Laurence L. Lesage-Meessen, et al.. Survey of fungal lignocellulose-acting enzymes for improved plant biomass degradation capacities. 2. Lund Symposium on Lignin and Hemicellulose Valorisation, Nov 2015, Lund, Sweden. 2015. hal-02792542

HAL Id: hal-02792542

<https://hal.inrae.fr/hal-02792542v1>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.


Survey of fungal lignocellulose-acting enzymes for improved plant biomass degradation capacities

Eric Record, Anne Lomascolo, Anne Favel, David Navarro, Laurence Lesage-Meessen, Annick Doan, Mireille Haon, Christophe Boyer, Sabine Taussac, Sacha Grisel, Emmanuel Bertrand, Sana Raouche, Isabelle Gimbert, Jean-Guy Berrin, Marie-Noëlle Rosso, Craig Faulds

INRA-AMU, UMR 1163 Fungal Biodiversity and Biotechnology, 163 avenue de Luminy CP 925 13288 Marseille Cedex 9, France (eric.record@univ-amu.fr)

Comparative genomics of fungal enzyme machinery

- Filamentous fungi are the most potent degraders of lignocellulosic biomass
- They secrete high number and a broad variety of carbohydrates-active enzymes and lignin-active enzymes (CAZymes) Levasseur et al. 2013. Lombard et al. 2013.




Genome sequencing and enzyme characterization

- CIRM a Biological Resource Center dedicated to the preservation of filamentous fungi of interest to agro-industries and their utilization (<http://www6.inra.fr/cirm/Champignons-Filamenteux>)
- The first genome sequence : *Pycnoporus cinnabarinus* : fungal model to study the enzyme mechanism to degrade lignin Levasseur et al. 2014.

Family	Total gene number
AA1 (multi-copper oxidases)	5 laccases <i>sensu stricto</i> , 1 Mco, 1 ferroxidase
AA2 (class II peroxidases)	12 (4 LiP, 1 VP, 1 atypical-VP, 3 short-MnP, 2 chloroP, and 1 partial protein)
AA3_1 (cellulose dehydrogenase)	1
AA3_2 (AAO, glucose oxidase/dehydrogenase)	19
AA3_3 (alcohol oxidase/dehydrogenase)	2
AA3_4 (pyranose oxidase)	2
AA5_1 (copper radical oxidases)	7 (3 Glyoxal oxidases <i>sensu stricto</i> , 4 Cro)
AA6 (benzoquinone reductase)	1
AA8	1
AA9 (LMPO)	15
Total	60

- Enzyme characterization, structure/function studies, chimerical enzymes (bi-functional), enzyme synergy
- Applications of enzymes for lignin valorization (Indox project)





Fungal secretome analysis

- Podospora anserina* : coprophilous fungus model adapted to complex biomass conversion
- Pestalotiopsis sp.* : mangrove fungal model to study salt-adapted enzymes

Poidevin et al. 2014


Podospora anserina




Wheat straw hydrolysis by *T. reesei* cellulases (E508) supplemented by *P. anserina* secretomes

Proteomic analyses

Avicel




Sugar Beet Pulp



Functional category analysis of the 142 proteins detected in *P. anserina* secretomes grown on Avicel and Sugar Beet Pulp.

Arfi et al. 2013.


Pestalotiopsis sp.



Rhizophora stylosa

Identification of salt tolerant enzymes : lignocellulolytic activity screening of isolated mangrove fungi after 8 days cultivation on non-saline or 3% saline conditions

Proteomic analyses



Salinity => Xylanases (GH43, GH10, GH30)
=> Cellulases (GH61, AA9 LPMO)

209 identified proteins with differential secretion : secretion of oxidoreductases found to be limited under saline conditions while cellulase and hemicellulase secretion were increased.