



HAL
open science

Tyrosine metabolism and parthenogenetic development in a symbiotic insect, the pea aphid: the central function of phenylalanine hydroxylase

Pierre Simonet, Karen Gaget, Marjolaine Rey, Marie-Gabrielle Duport, Hubert Charles, Gérard Febvay, Stefano Colella, Federica Calevro

► To cite this version:

Pierre Simonet, Karen Gaget, Marjolaine Rey, Marie-Gabrielle Duport, Hubert Charles, et al.. Tyrosine metabolism and parthenogenetic development in a symbiotic insect, the pea aphid: the central function of phenylalanine hydroxylase. 7. International Symposium on Molecular Insect Science, Jul 2014, Amsterdam, Netherlands. 1 p., 2014. hal-02792558

HAL Id: hal-02792558

<https://hal.inrae.fr/hal-02792558>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Title:

Tyrosine metabolism and parthenogenetic development in a symbiotic insect, the pea aphid: the central function of phenylalanine hydroxylase

Authors & affiliations:

P. Simonet, K. Gaget, M. Rey, G. Duport, H. Charles, G. Febvay, S. Colella, F. Calevro
UMR203 BF2I, INRA, INSA-Lyon, Université de Lyon, France
federica.calevro@insa-lyon.fr

Abstract:

Nutritional symbioses are widespread associations among insects, contributing to the evolutionary adaptation and the invasive power of this class of animals. The endosymbiosis between the pea aphid, *Acyrtosiphon pisum*, and the bacterium, *Buchnera aphidicola*, is an interesting model of symbiotic partner's integrated metabolism, particularly for the production of amino acids lacking in the aphid's diet. An important supply of these nutrients has to be provided especially for the phenomenal aphid growing rate observed during parthenogenetic viviparous reproduction.

A recent transcriptomic analysis, performed in our laboratory, allowed us to identify a key metabolic pathway regulated in the pea aphid parthenogenetic development: the tyrosine pathway. This biosynthetic pathway is shared by the two symbiotic partners, all the precursors being produced by *B. aphidicola*. To determine the involvement of this pathway in aphid development, we silenced by RNAi approach the unique gene in the pea aphid genome that encodes for phenylalanine hydroxylase (*ACYPI007803*, EC 1.14.16.1), the specific enzyme of tyrosine biosynthesis catalyzing the conversion of Phe to Tyr.

ACYPI007803 knockdown by microinjection of dsRNA yielded several defective phenotypes, specifically targeting the two symbiotic compartments: the bacteriocytes and the embryo chains. The *ACYPI007803* gene expression inhibition, confirmed by qRT-PCR, considerably affected aphid fecundity by diminishing the number of nymphs produced. Furthermore, the RNAi-mediated treatment resulted in a significant impairment of embryo development, with the production of non-viable and deformed neonatal nymphs. To attest of the tyrosine biosynthesis disruption following *ACYPI007803* knockdown, single embryo HPLC analyses were performed and demonstrated a significant reduction of tyrosine amount as well as a coherent increase of phenylalanine, the upstream amino acid precursor.

Altogether, our findings shed new light on the importance of the tyrosine pathway in the embryonic development of parthenogenetic pea aphids and establish the phenylalanine hydroxylase contribution to this process, notably by regulating the tyrosine biosynthesis.