



HAL
open science

Identification of candidate causal variants underlying QTL in dairy cattle through GWAS and Bayesian approach at the sequence level

Didier Boichard, Marie-Pierre Sanchez, Anne Barbat, Mekki Boussaha, Thierry Tribout, Rachel Lefebvre, Sebastien S. Fritz, Romain Saintilan, Chris Hoze Hoze, Armelle Gion Govignon-Gion, et al.

► To cite this version:

Didier Boichard, Marie-Pierre Sanchez, Anne Barbat, Mekki Boussaha, Thierry Tribout, et al.. Identification of candidate causal variants underlying QTL in dairy cattle through GWAS and Bayesian approach at the sequence level. PAG XXIV - Plant and Animal Genome Conference, Jan 2016, San Diego, United States. hal-02792675

HAL Id: hal-02792675

<https://hal.inrae.fr/hal-02792675v1>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



PLANT & ANIMAL GENOME CONFERENCE XXIV

The Largest Ag-Genomics Meeting in the World.

January 9-13, 2016
San Diego, CA
www.intlpag.org

W424

Identification of Candidate Causal Variants Underlying QTL in Dairy Cattle through GWAS and Bayesian Approach at the Sequence Level

Date: Saturday, January 9, 2016

Time: 4:05 PM

Room: Golden West

Didier A. Boichard , INRA UMR1313 Animal Genetics and Integrative Biology, Jouy en Josas, France
Marie-Pierre Sanchez , INRA UMR1313 Animal Genetics and Integrative Biology, Jouy en Josas, France
Anne Barbat , INRA UMR1313 Animal Genetics and Integrative Biology, Jouy en Josas, France
Mekki Boussaha , INRA UMR1313 Animal Genetics and Integrative Biology, Jouy en Josas, France
Thierry Tribout , INRA UMR1313 Animal Genetics and Integrative Biology, Jouy en Josas, France
Rachel Lefebvre , INRA UMR1313 Animal Genetics and Integrative Biology, Jouy en Josas, France
Sébastien Fritz , Alice, Genetics Team, Paris, France
Romain Saintilan , Alice, Genetics Team, Paris, France
Chris Hoze , Alice, Genetics Team, Paris, France
Armelle Govignon-Gion , INRA UMR1313 Animal Genetics and Integrative Biology, Jouy en Josas, France
Pascal Croiseau , INRA UMR1313 Animal Genetics and Integrative Biology, Jouy en Josas, France

Combining large genotyped and phenotyped resource populations with whole genome sequence information opens new avenues to unravel candidate mutations underlying the genetic variability of the phenotypic traits. Reference populations for genomic selection are very large (more than 100,000 dairy animals with phenotypes and genotypes in France) and increase rapidly. As shown by the 1000 bull genomes project (1682 sequenced bulls in 2015), the number of sequenced bovine genomes also increases very fast and already makes it possible to impute the reference populations up to the sequence level, although rare alleles still have limited imputation accuracy. These populations sequenced in silico are unique resources for GWAS studies. Several strategies can be applied: first, each population can be analyzed by simple GWAS one marker at a time; results for the same trait but from multiple breeds can be combined in meta-analyses; multi-markers approaches can be implemented on targeted regions (1-3 Mb) in order to reduce the effect of long range linkage disequilibrium; haplotype analyses can be used to filter out genetic variants with inconsistent effects; finally, functional annotations help selecting the best candidates and interpreting their likely effects. Examples from the PhenoFinLait project are given, showing that QTL involved in milk protein or fatty acids composition can be characterized with a high confidence up to the gene and, frequently, up to a very limited number of candidate mutations. The authors acknowledge the financial support from ANR and APIS-GENE, and the contribution of the 1000 bull genomes consortium.

Back to: [Genomic Selection and Genome-Wide Association Studies](#)

[Previous Abstract](#) | [Next Abstract >>](#)

[Home/Search](#)

[Browse by Day](#)

[Browse by Type](#)

[Poster Categories](#)