

Design of a SNP parentage assignment panel for French goat breeds

Isabelle Palhiere Palhière, Flavie Tortereau, Pierre Martin, Rachel Rupp,

Gwenola Tosser-Klopp

► To cite this version:

Isabelle Palhiere Palhière, Flavie Tortereau, Pierre Martin, Rachel Rupp, Gwenola Tosser-Klopp. Design of a SNP parentage assignment panel for French goat breeds. XXIV. Plant and Animal Genome Conference (PAG), Jan 2016, San Diego, United States. 1 p. hal-02793374

HAL Id: hal-02793374 https://hal.inrae.fr/hal-02793374v1

Submitted on 5 Jun2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Design of a SNP parentage assignment panel for French goat breeds

Palhière I.(1), Tortereau F.(1), Martin P.(2), Rupp R.(1), Tosser-Klopp G.(1)
(1) INRA, INPT ENSAT, INPT ENVT, UMR 1388 Génétique, Physiologie et Systèmes
d'Elevage, F-31326 Castanet-Tolosan, France
(2) Capgènes, F-86550 Mignaloux-Beauvoir, France

Knowing pedigrees is essential for selection, conservation and management of animal populations. In French goats, complete and accurate pedigree information is difficult to get because of the increasing use of multiple-sired matings. In the two main French breeds (Alpine and Saanen), only 50% of females involved in the official genetic evaluation have a known sire. In this context, having an efficient way to *a posteriori* identify actual parentage could be profitable both at the breeding scheme and the breeder levels. The objective of this study was to develop a SNP panel to accurately assign paternity in the main French goat breeds. About thirty animals have been genotyped with the GoatSNP50 chip in eight breeds: Alpine, Saanen, Angora, Corsican, Poitevine, Pyrénées, Fossés, Provençale. Four subsets of SNP (64 to 505) were theoretically defined based on call frequency, mendelian incompatibility, Hardy-Weinberg equilibrium test, Minor Allele Frequency and distribution along the genome. Then, the assignment rate of each SNP panel was estimated on Alpine and Saanen data from a QTL detection design, and by using a likelihood based approach. The 505 and 246 SNP panels showed the best results: the average MAF was about 0.42 for all the markers and populations, and the assignment rate was 100%. Before choosing the final panel, an assignment test will be performed in farm conditions (collection of blood samples, multiple-sired matings,...). A selection of SNP located in major genes of interest will be added to those for assignment in order to provide a useful tool to industry.