

Spatial pattern of Eurasian woolly mammoth extinctions during the Late Pleistocene based on statistical inference from fossil records

Frédéric Saltré, Joel Chadoeuf, B.W. Brook, Damien A. Fordham, Marta Rodriguez-Rey, Corey J.A. Bradshaw

► To cite this version:

Frédéric Saltré, Joel Chadoeuf, B.W. Brook, Damien A. Fordham, Marta Rodriguez-Rey, et al.. Spatial pattern of Eurasian woolly mammoth extinctions during the Late Pleistocene based on statistical inference from fossil records. 9. IALE World Congress, International Association of Landscape Ecology (IALE). USA., Jul 2015, Portland Oregon, U.S. Virgin Islands. hal-02793379

HAL Id: hal-02793379

<https://hal.inrae.fr/hal-02793379>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Saltré, F., Chadoeuf, J., Brook, B.W., Fordham, D.A., Rodriguez-Rey, M., & Bradshaw, C.J.A., (2015). Spatial pattern of Eurasian woolly mammoth extinctions during the Late Pleistocene based on statistical inference from fossil records. 9th IALE World Congress, Portland, OR, USA.

Title

Spatial pattern of Eurasian woolly mammoth extinctions during the Late Pleistocene based on statistical inference from fossil records

Authors

F. Saltré, B.W. Brook, J. Chadoeuf, D.A. Fordham, M. Rodriguez-Rey, C.J.A. Bradshaw

Abstract:

Late Pleistocene extinctions affected most large and low-fecundity mammals such as the woolly mammoth (*Mammuthus primigenius*) in Eurasia. Despite recent advances in both genetic and modelling approaches and increasing fossil data, assessing the relative importance of climate variations and human hunting on the spatial pattern of mammoth extinction remains a challenge. This is because estimating incomplete fossil time series biases the true dates of extinction and appearance, potentially leading to incorrect conclusion of the relative importance of individual and synergistic drivers. We compared outputs of two statistical models that (i) spatially infer mammoth final extinction dates and first human occurrence from sighting records based on the recently developed Gaussian-resampled inverse-weighted McInerny (GRIWM) and (ii) predict the date when mammoths should have gone extinct based on climate change only (climate anomaly and velocity). GRIWM generates a confidence interval of the true extinction time, which suggests a pattern of mammoths extinctions starting in central Europe and spreading toward both western Europe and north-eastern Siberia. Climate change had a minor impact on mammoth extinctions, mainly in the northeastern part of Eurasia where the synergistic hypothesis is mainly supported by our results whereas the human hunting hypothesis is more likely to have caused mammoth extinctions elsewhere in Eurasia.