



HAL
open science

Quantitative in-situ NMR to characterize protein oxidation and its dynamics

Guilhem Pages, Amaël Morisse, Philippe P. Gatellier, Estelle Martineau,
Patrick Giraudeau, J.-M. Bonny

► To cite this version:

Guilhem Pages, Amaël Morisse, Philippe P. Gatellier, Estelle Martineau, Patrick Giraudeau, et al.. Quantitative in-situ NMR to characterize protein oxidation and its dynamics. 13. International Conference on the Applications of Magnetic Resonance in Food Science, Jun 2016, Karlsruhe, Germany. 1 p. hal-02793518

HAL Id: hal-02793518

<https://hal.inrae.fr/hal-02793518>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Quantitative *in-situ* NMR to characterize protein oxidation and its dynamics

G. Pagès¹, A. Morisse^{1,2}, P. Gatellier¹, E. Martineau^{2,3}, P. Girardeau^{2,4}, J.-M. Bonny¹

¹AgroResonance, UR370 QuaPA - INRA, F-63122 Saint-Genès-Champanelle, France

²EBSI Team, Chimie et Interdisciplinarité : Synthèse, Analyse, Modélisation (CEISAM), Université de Nantes, CNRS, UMR 6230, LUNAM Université, France

³Spectromaîtrise, CAPACITES SAS, Nantes, France

⁴Institut Universitaire de France, 1 rue Descartes, 75005 Paris Cedex 5, France

Preserving food quality is critical to limit the oxidation processes. The evolution of meat colour or the development of rancid taste in oils are two examples of oxidative processes degrading the food quality. The reaction of oxygen (or its derivatives) with metal ions naturally present in food (*eg.* iron) forms free radical reactive oxygen species (ROS). These ROS are the main factors of food oxidation. The aim of this work is to evaluate the intakes of quantitative *in situ* NMR to understand and characterize the oxidation mechanisms.

Our preliminary work focussed on the evaluation of some amino acid mixtures as models of protein oxidation. Due to NMR signal overlaps, recording 2D NMR spectra is indispensable to isolate NMR signals from targeted amino-acids. However, these experiments are time-consuming and not adapted to chemically evolving media. To address this limitation, we developed tailored hybrid methods based on ultrafast 2D NMR. The spectrum recording time decreased from ~30 min for a classical pulse sequence to a few minutes only with the ultrafast method. This approach allows the real-time monitoring of chemical evolutions in such complex mixtures. Using this quantitative approach, we observed a fast oxidation for the histidine while threonine and lysine oxidization kinetics were significantly slower. Our analytical approach offers a promising tool to monitor oxidation processes in food products.