



HAL
open science

Transfer of a miniaturized method for high-throughput screening of biomass pretreatment and saccharification and application on poplar and miscanthus clones

Nassim Belmokhtar, Vincent Segura, Brigitte Chabbert, Nathalie Boizot, Kévin Ader, Stéphanie Arnoult, Maryse Brancourt Brancourt-Hulmel, Jean-Charles Bastien, Catherine Bastien, Jean-Paul J.-P. Charpentier

► To cite this version:

Nassim Belmokhtar, Vincent Segura, Brigitte Chabbert, Nathalie Boizot, Kévin Ader, et al.. Transfer of a miniaturized method for high-throughput screening of biomass pretreatment and saccharification and application on poplar and miscanthus clones. Exploring lignocellulosic biomass! Explorons la biomasse lignocellulosique!, Jun 2016, Reims, France. , 2016. hal-02793860

HAL Id: hal-02793860

<https://hal.inrae.fr/hal-02793860>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Transfer of a miniaturized method for high-throughput screening of biomass pretreatment and saccharification and application on poplar and miscanthus clones


Nassim Belmokhtara,b, Vincent Seguraa, Brigitte Chabbertc, Nathalie Boizota,b, Kévin Adera,b, Stéphanie Arnouldt, Maryse Brancourt-Hulmele, Jean-Charles Bastiena, Catherine Bastiena, **Jean Paul Charpentiera,b**

a INRA, Centre Val de Loire, UR 0588 AGPF, 45075 Orléans Cedex 2, France
 bINRA, Centre Val de Loire, Plateforme Régionale GéoBois, 45075 Orléans Cedex 2, France
 cINRA, Centre Nord Picardie Champagne, UMR 0614 FARE, 51100 Reims, France
 dINRA, Centre Nord Picardie Champagne, UE GCIE, Estrée-Mons, 80203 Péronne, France
 eINRA, Centre Nord Picardie Champagne, UR AgrolImpact, Estrée-Mons, 80203 Péronne, France

Introduction

In order to meet increasing demand for second-generation biofuels, exploring large sets of lignocellulosic biomasses appears to be one of the key levers to improve bio refinery yields.


Poplar and miscanthus breeding programs have provided large sets of varietal outputs, whose saccharification potential assessment could lead to identify the best biofuel-oriented varieties.



Objectives of this study:


- 1- Transfer into breeding programs and adaptation of a high-throughput screening method developed at NREL (USA)¹
- 2- Demonstration, on poplar and miscanthus biomass, of the importance of pretreatment step in screening process

Experiments




Results

1- Impact of pretreatment on the variability of glucose release percentage



- Hot water pretreatment in 96-well microplates increases significantly saccharification yield for both plants
- More variability in glucose release is observed after hot water pretreatment mainly for poplar clones

2- High-throughput screening of 10 SRC and 10 vSRC poplar varieties



- vSRC poplars release higher glucose amounts than SRC ones when saccharified without pretreatment (A)
- SRC and vSRC poplars show similar enzymatic glucose yield after hot water pretreatment (B)
- More variability in glucose release is observed for SRC poplar without pretreatment (A) in contrast to vSRC ones which express more variability after hot pretreatment (B)

Conclusions

- Based on the comparison of the sugar release of poplar and miscanthus biomasses, our approach allowed us to stress the importance of a pretreatment step before enzymatic hydrolysis to magnify the expression of the genetic variability. Furthermore, this approach revealed different responses to saccharification of SRC and vSRC poplar biomasses without or after a pretreatment.
- Our developed high-throughput method is well adapted to screen, with high accuracy, a large set of samples from different lignocellulosic biomasses to evaluate their potential for saccharification.

Reference:

[1] M.J. Selig, M.P. Tucker, R.W. Sykes, K.L. Reichel, R. Brunecky, M.E. Himmel, M.F. Davis and S. R. Decker. Lignocellulose recalcitrance screening by integrated high-throughput hydrothermal pretreatment and enzymatic saccharification. *Industrial Biotechnology* 6 (2010), 104-111.

Acknowledgments:

This work was funded by BPIFrance in the R&D project Futurol
 Innovative equipments were funded by Xyloforest, an Equipex Project (ANR) (INRA Aquitaine)

E-mail contacts :

Nassim.Belmokhtar@orleans.inra.fr
 Jean-Paul.Charpentier@orleans.inra.fr