



Effect of the spatial distribution of resources in experimental mini-landscapes on the fitness of two populations of *Trichogramma cacoeciae*

Victor Burte, Guy Perez, Ludovic Mailleret, Vincent Calcagno

► To cite this version:

Victor Burte, Guy Perez, Ludovic Mailleret, Vincent Calcagno. Effect of the spatial distribution of resources in experimental mini-landscapes on the fitness of two populations of *Trichogramma cacoeciae*. International Conference on Ecological Sciences, Oct 2016, Marseille, France. hal-02794095

HAL Id: hal-02794095

<https://hal.inrae.fr/hal-02794095>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effect of the spatial distribution of resources in experimental minilandscapes on the fitness of two populations of *Trichogramma cacoeciae*.

Victor Burte^{*1}, Guy Perez¹, Ludovic Mailleret¹, and Vincent Calcagno¹

¹Institut Sophia Agrobiotech [Sophia Antipolis] (ISA) – Institut national de la recherche agronomique (INRA) : UMR1355, Université Nice Sophia Antipolis (UNS) – INRA Centre de recherche Provence-Alpes-Côte d’Azur 400, route des Chappes BP 167 06903 Sophia Antipolis Cedex, France

Abstract

Parasitoids in genus *Trichogramma* (Hymenoptera) are minute size (less than 1 mm) wasps that lay eggs in the eggs of other insect species. They have been used in biological control for decades to contain Lepidoptera pest populations in many different crops. They may adapt their behaviour to maximize their performance at finding their host eggs, which are a patchy and heterogeneously distributed resource. To understand how *Trichogramma* are affected by the spatial distribution of their resource, we assembled experimental minilandscapes in two dimensions. We created different resource distributions varying the degree of spatial autocorrelation (aggregation) of *Ephesttia kunhiella* eggs, keeping average egg density constant. We studied two populations of *Trichogramma cacoeciae*, the first (PJ) maintained under laboratory conditions for more than twenty years and the second (WT) sampled in the field in 2014. For each population we determined parasitism performance (fitness) and monitored individual movements over the landscapes using high-resolution pictures over the entire foraging sequences (five hours). We found that resource spatial distribution had a strong impact on fitness for both populations, but that the two populations responded in different ways: PJ was more efficient than WT when egg distribution was either very homogeneous or very heterogeneous (aggregated), whereas WT was more efficient than PJ at intermediate levels of spatial autocorrelation. We will show how these fitness differences can be explained from differences in the underlying individual behaviours and movement patterns.

Keywords: optimal foraging, mini landscape, trichogramma, patch, picture analysis

^{*}Speaker