



Biotic and abiotic factors influence the expression of effectors in *Leptosphaeria maculans* during axenic growth

Michel M. Meyer, Ahmed Salim A. S. Bourras, Clémence Plissonneau, Julie Gervais, Marie-Helene Balesdent, Thierry T. Rouxel

► To cite this version:

Michel M. Meyer, Ahmed Salim A. S. Bourras, Clémence Plissonneau, Julie Gervais, Marie-Helene Balesdent, et al.. Biotic and abiotic factors influence the expression of effectors in *Leptosphaeria maculans* during axenic growth. ECFG13. European conference on fungal genetics (2016), Apr 2016, PARIS LA VILLETTE, France. p.309, 2016. hal-02794169

HAL Id: hal-02794169

<https://hal.inrae.fr/hal-02794169>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER SESSION ABSTRACTS
Session CS3 Environmental sensing, stress response
CS3T25

Tuesday 5th April
14:00 - 16:00

MEYER Michel (1), BOURRAS Salim (2), PLISSONNEAU Clémence (1), GERVAIS Julie (1), BALESDENT Marie-Hélène (1), ROUXEL Thierry (1)

(1) BIOGER, INRA, AgroParisTech, Thiverval-Grignon, France

(2) Institute of Plant Biology, University of Zürich, Switzerland

Biotic and abiotic factors influence the expression of effectors in *Leptosphaeria maculans* during axenic growth

Plant pathogens secrete effector proteins into host tissues to promote infection through the manipulation of host processes. Sequencing and analyses of the genomes of fungal phytopathogens have shown that they contain tens to hundreds of genes predicted to encode putative effectors. Moreover, global analyses of gene expression revealed that several waves of concerted expression of effector genes take place during host invasion. In sharp contrast with the situation described in planta, the expression of the effectors is difficult to detect and quantify in axenic cultures because their genes are expressed at a very low level. In the present study, we investigate biotic and abiotic factors that may relieve suppression of expression of effectors during axenic growth. Biotic factors (such as carbon source, nitrate source, antibiotics) as well as abiotic factors (pH, temperature) can influence their expression. Of major interest, incubation of the fungal mycelium with 1 µg/ml of an antibiotic of the aminoglycoside family allowed an increase of effector gene expression 20-fold to 60-fold compared to regular axenic growth. An RNAseq analysis aiming at identifying the set of effectors up- and down-regulated in a culture medium supplemented or not with the antibiotic has been performed and results obtained will be presented. This simple system could be a good starting point to characterize the plant signals that trigger fungal effector gene expression.
