Root development of non-accumulating and hyperaccumulating plants in metal contaminated soils amended with biochar

Frédéric Rees, Thibault Sterckeman, Jean-Louis Morel

► To cite this version:

Frédéric Rees, Thibault Sterckeman, Jean-Louis Morel. Root development of non-accumulating and hyperaccumulating plants in metal contaminated soils amended with biochar. 8. 20th World Congress of Soil Science, Jun 2014, Jeju, North Korea. 2014. hal-02794196

HAL Id: hal-02794196

<https://hal.inrae.fr/hal-02794196>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

ROOT DEVELOPMENT IN METAL CONTAMINATED SOILS AMENDED WITH BIOCHAR

Frédéric REES¹, Thibault STERCKEMAN¹, Jean-Louis MOREL¹

¹Université de Lorraine/INRA, Laboratoire Sols et Environnement
F-54500 Vandœuvre-lès-Nancy, France

*corresponding author: Frédéric REES, f.rees@gisfi.fr

INTRODUCTION

Biochar, the solid product from biomass pyrolysis used as soil amendment, has emerged as a promising carbon sink and soil improver. Its sorbent properties could also be used in the remediation of contaminated soils, particularly in **phytoremediation**.

Biochar's influence on **root growth** is however poorly known ^[1], e.g. for **soils contaminated with heavy metals**. An increase of root surface in those soils may lead to a **decrease of metal leaching**, as less water is percolating, but also to an **increase of metal uptake** by the plant, as the exchange surface between soil and plant is increasing.

In this context, several mechanisms ^[1,2,3] could explain a better root development:

- [1] Soil **toxicity** ↘
- [2] **Water** availability ↗
- [3] Resistance to **root penetration** ↘
- [4] **Nutrient** availability ↗
- [5] Beneficial **microorganisms** are promoted
- [6] Biochar induces plant **hormonal response**

→ A **rhizobox** experiment was designed to:

- 1) Quantify the effect of biochar on **root growth in contaminated soils**
- 2) Identify a possible **root tropism** towards biochar, thanks to a specific design

MATERIALS

- **1 biochar** produced by Carbon Terra at ~450°C from woody biomass, <2mm, untreated (pH 9.2)
- **2 soils** contaminated with Cd, Zn, Pb, sampled near smelters, with similar properties but different pH.

pH value and available (/ total) metals of soils (mg kg⁻¹)

Soil	pH	Cd	Pb	Zn
Soil A	5.9	5.9 / 17.6	1.7 / 1120	684 / 3170
Soil B	8.1	0.24 / 18.6	0.06 / 1080	2.0 / 1380

→ With biochar, metal availability strongly ↘ on Soil A, but only slightly on Soil B due to its higher initial pH ^[4]

- **2 plant species** grown in large rhizoboxes:
 - Zea mays, non hyperaccumulating, fast growing
 - Alpine pennycress, Cd and Zn hyperaccumulator

METHODS

- 4 seedlings grown per rhizobox with 2000 g of soil, divided in 8 compartments as a **chessboard**:

4 squares with pure soil
4 squares with soil + 5% (w/w) biochar

- 2 rhizoboxes for each plant and soil
- Daily watering at 85% of water holding capacity
- High-resolution scanning of the soil profile

▪ After harvest (2 weeks for maize, 9 weeks for alpine pennycress):
→ Recovery of roots and measurement of root surface with Winrhizo software

DEVELOPMENT OF ZEA MAYS ROOTS

22	24	22	30
73	20	58	9

RESULTS & DISCUSSION

← ON SOIL A →

- The zones with biochar have a **higher density of roots**
- Roots are generally moving towards the zones with biochar
→ **root tropism**

=> Biochar has **clear positive effects** on root development in a soil with **initial high metal availability and low pH**

← ON SOIL B →

- The zones with biochar do not have a higher density of roots
- No obvious trend of root tropism towards biochar can be observed

=> Biochar has **no significant effects** on root development in a soil with **initial low metal availability and high pH**

DEVELOPMENT OF ALPINE PENNYCRESS ROOTS

16	42	25	51
174	71	141	79

10	11	11	18
29	19	23	29

Tables represent the total root surface for the 8 squares of each soil profile (in cm²)

CONCLUSIONS

- Considering that both soils have similar properties except pH, the **better root development with biochar** only observed on Soil A may be mainly **due to the decrease of soil metal availability**.

→ Modifications of root development only occurs when biochar has a significant effect on chemical soil properties.

PERSPECTIVES

- Positive tropism of roots towards biochar could be an option to reduce the quantity of biochar and the work for biochar amendment.
- The consequences of a better root development on plant metal uptake and long term growth need further investigations.

[1] Prendergast-Miller, M.T., Duvall, M., Sohi, S.P. 2013. Biochar-root interactions are mediated by biochar nutrient content and impacts on soil nutrient availability. European Journal of Soil Science.

[2] Jones et al. 2012. Biochar-mediated changes in soil quality and plant growth in a three year field trial. Soil Biology and Biochemistry.

[3] Spokas, K.A., Baker, J.M. and Reicosky, D.C. 2010. Ethylene: potential key for biochar amendment impacts. Plant and Soil.

[4] Rees et al. 2014. Short-term effects of biochar on soil heavy metal mobility are controlled by intra-particle diffusion and soil pH increase. European Journal of Soil Science.