



**HAL**  
open science

## High-resolution spatial modelling of total soil depth for France

Marine Lacoste, Vera Leatitia Mulder, Nicolas N. Saby, Dominique D.  
Arrouays

► **To cite this version:**

Marine Lacoste, Vera Leatitia Mulder, Nicolas N. Saby, Dominique D. Arrouays. High-resolution spatial modelling of total soil depth for France. 6. Global Workshop on Digital Soil Mapping, Nov 2014, Nanjing, China. 11 p. hal-02794346

**HAL Id: hal-02794346**

**<https://hal.inrae.fr/hal-02794346v1>**

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# High-resolution spatial modelling of total soil depth for France

**Marine Lacoste** – INRA UR 0272 Science du Sol, Orléans, France

**Titia Mulder** – INRA US Infosol, Orléans, France

Contributors: M. Martin, N. Saby, A. Richer de Forges & D. Arrouays

Project: GlobalSoilMap

6th Global Workshop on Digital Soil Mapping

11-14 November, 2014, Nanjing, China



# INTRODUCTION

## ❖ Soil depth (SDt):

- Key soil property for water availability and carbon stocks
- Exhaustive mapping of total soil depth = requirement of the GlobalSoilMap project

## ❖ Difficulties of SDt mapping due to:

- Soil properties: high spatial variability
- Soil observation tools: estimation of soil depth for deep soils (> 1.5 m)
- Discordance about SDt definition

## ➤ Evaluate two different modelling approaches to produce a high-resolution soil depth map of France

- In a regional or global context + high resolution
  - Large data sets
  - Spatial heterogeneity
  - Local, large and nested-scale processes
- Robust and reproducible
- Spatial explicit uncertainties



# RESEARCH OVERVIEW

## Input data

- **Soil sample data**  
(source: French Soil Monitoring network)
- **Exhaustive covariates** capturing biotic and abiotic conditions
  - Soil type and properties
  - Parent material
  - Relief (SRTM-DEM)
  - Climate
  - Land use

## Analysis

### 1) Data mining

- + Bias correction
- + Ordinary kriging of the residuals

*Resolution: 90 m*

R packages: *caret*, *gbm*,  
*qmap*, *gstat*

### 2) Multi-resolution kriging for large datasets

*Fixed trend model + kriging*

*Resolution: 500 m*

R packages: *LatticeKrig*

## Evaluation criteria


- 1) Map accuracy
  - Internal validation
  - Cross-validation
  - External validation:  
concordance with previous soil map
- 2) Prediction and confidence intervals by conditional simulation of kriging model

# RESULTS (I/IV)

Data mining

Multi-resolution

Validation Type (90% prediction interval)	Data mining	Multi-resolution Kriging
Internal	91 %	32 %
External	72 %	30 %



	Min	Q1	Mean	Median	Q3	Max	sd
Data Mining	0	97	111	113	127		197
MR Kriging	13	35	38	38	42	72	6
Difference	-154	-18	4	1	24	186	33

As discussed: the validation is incorrect for MR Kriging

# Discussion

## Data mining

### Predictive map of soil depth

Consistent spatial pattern

Good prediction of the mean values

- Prediction of extremes values

**90% Confidence interval**

- **Large (high uncertainties)**
- “Consistent” with observed values

### Implementation

- **Multisteps/multitools approach**
- **No direct estimation of uncertainties**
- Flexible for large datasets, high resolution

- Promising prediction of soil depth class instead

## Multi-resolution Kriging

Ongoing: increasing the resolution/levels

- Narrow (low uncertainties)

Ongoing: test lower confidence intervals

- Straight forward modelling approach
- Flexible in delivering spatial explicit uncertainty measures

### Outlook

- Potential for modelling beyond the country level, at high resolution as demonstrated in other global environmental models



# THANK YOU ALL!

*Essentially, all life depends upon the soil.  
There can be no life without soil and no soil without life;  
they have evolved together.*

American naturalist Charles Kellogg, 1938.

## FINANCIAL SUPPORT:




Inventaire, gestion  
et conservation  
des sols



Région Centre


# STUDY AREA : France (~ 540K km<sup>2</sup>)

16km x 16km grid



SDt determined for 2116 sites

French Soil Monitoring network (RMQS)



Mean value: 102  
cm




# STUDY AREA : France (~ 540K km<sup>2</sup>)

Existing soil depth maps

Scale: 1/1 000 000

Lower limit


Upper limit



Scale: 1/250 000

Lower limit

Upper limit



Note: these are classes. The spatial distribution of these classes is the same as the soil type classes used for the data mining model: this introduces bias in the following validation results – I have my questions about that approach...

# METHODS

## Continue soil depth prediction

### Data mining

### Multi-resolution Kriging

- Estimation of covariance matrix using multi-resolution radial basis functions
- Covariance model can approximate the Matern covariance family
- Developed for handling large datasets
- *R package LatticeKrig*
- Resolution: 500m – for me it doesn't make sense to go to 90m because it is not supported by the data we use....also, the model cannot be calibrated because there is no variability below this level
- Fixed linear trend model: elevation, slope, precipitation, gravimetry, bed rock resistance and NPP
- **Kriging error obtained by conditional Gaussian simulation (1000 times) – this is really a pro!**

# RESULTS

## Importance of the covariates

### Data mining

Variable	Importance (%)
SRTM (elevation)	14
Maximal annual temperature (mean)	9
Parent material	8
Aspect	7
Mean annual precipitation	7
Climate type	7
Roughness	7
Land use for forest areas	6
Wetness index	6
Soil type	6
Drainage network	6
Slope position	6
Slope	6
Bare rock areas	5

### Multi-resolution Kriging

- Fixed linear trend model: elevation, slope, precipitation, gravimetry, bed rock resistance and NPP – what are the coefficients?

A vertical photograph of a soil profile. On the left, a black measuring tape is visible, showing depth in centimeters from 0 to 120. The soil layers are visible, with a dark top layer, a lighter brown layer, and a darker, more textured layer at the bottom.

# RESULTS

## Models accuracy

Interesting to see the multi-resolution kriging improves with a higher resolution soil class map. The good validation results for data mining relate to the previous mentioned bias. The classes have been very important for the data mining – this data does not have the spatial variability compared to eg SRTM. Matching the soil depth class with modelled soil depth thus shows high agreement + The variogram of the residuals did not show high spatial variability. Concluding – a soil class map is not the best type of validation here. BTW the internal validation of the MR kriging is cross validation – so not too bad compared to the cross-validation of the data mining technique.

The histograms should be changed to relative frequency due to the different resolution – or make 2 separate histograms (difference in resolution = different total). The kriging, as expected, shows a smoothing of values (no extremes). What about the validation with the independent IGCS soil depth data? Still impossible because of the inaccuracy of that dataset? Maybe Anne knows how to select the most accurate samples – perhaps a specific year, institute or sampling program which was consistent over the years?