



HAL
open science

Improvement of vineyard sustainability according to biogeochemical cycle of nitrogen in field

Marie Thiollet-Scholtus, Najat Nassr, Aude Langenfeld, Mohamed Benbrahim, Lionel Ley, Julie Grignon, Christian C. Bockstaller

► To cite this version:

Marie Thiollet-Scholtus, Najat Nassr, Aude Langenfeld, Mohamed Benbrahim, Lionel Ley, et al.. Improvement of vineyard sustainability according to biogeochemical cycle of nitrogen in field. The 5th International Symposium for Farming Systems Design, Sep 2015, Montpellier, France. 2 p. hal-02794498

HAL Id: hal-02794498

<https://hal.inrae.fr/hal-02794498v1>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Improvement of vineyard sustainability according to biogeochemical cycle of nitrogen in field

Marie Thiollet-Scholtus ^{±1}, Najat Nassr ², Aude Langenfeld ², Mohamed Benbrahim ², Lionel Ley ³, Julie Grignion ⁴ & Christian Bockstaller ⁴

¹ INRA-SAD-UR-0055 ASTER, 28, rue de Herrlisheim F-68021 Colmar, France

² CRITT-RITTMO-Agroenvironnement, 37 Rue de Herrlisheim, 68000 Colmar, France

³ INRA-SEAV, 28, rue de Herrlisheim F-68021 Colmar, France

⁴ INRA-UMR-1121 LAE, 28, rue de Herrlisheim F-68021 Colmar, France

* Speaker

[±] Corresponding author: marie.thiollet-scholtus@colmar.inra.fr

1 Introduction

Good and yield wine production implies a well-balanced biogeochemical cycle of nitrogen (BCN) at field level e.i. in soil and in plant (Guilpart, *et al.*, 2014). Nitrogen is very important for grape quality and quantity and field sustainability. The mineralization of organic nitrogen, depending on soil microbial activity which is linked to soil cover crop management, is the main source of mineral nitrogen for the vine (Barlow *et al.*, 2009; Ingels *et al.*, 2005; Raath and Saayman, 1995; Thiebeau *et al.*, 2005). This paper is focused on a useful indicator fulfilling a sustainability assessment method: functional microbial populations implicated in BCN in vineyard field.

2 Materials and Methods

An experimental network with 6 platforms gathering 45 fields located in Atlantic coast (Loire valley and Bordeaux) and in North-East (Alsace) of France has been set up since 2012 (extract presented in Table 1). These vine sites represent a diversity of environmental factors (i.e. soil and climate) and the same method is used to assess them: agricultural, environmental, social and economical. The tested prototypes are tested according to the following goals: same yield, harvest quality, working time and production costs. The added value approach is based on assessing nitrogen dynamic in soil, i.e. nitrogen mineralization, regarding microbial biomass and activity.

Table 1. Alsacian sites-system characteristics.

Name	Site	Designed system	Vine age	Variety	Soil
Ribeau_AB	INRA-Ribeauvillé	Organic	16	Riesling	Loamy, sandy and clayey
Ribeau_PI		Integrated	16		
Rouff_PI	Rouffach agriculture school	Integrated	32	Pinot Gris	
Rouff_Piopti		Integrated-Rate-sprayingreduction	32		
Chaten_AB	OPABA-Chatenois	Organic and Biodynamic	17	Riesling	Sandy silty and clayey
Inger_AB	OPABA-Ingersheim		26		Sandy loam

3 Results – Discussion

For all AB-systems, functional richness (Fig.1A) at bud-break is very much higher than for all PI-systems (respectively 24-25 and 16-19). For sites Ribeauvillé and Rouffach, functional richness is not significantly different between designed systems at bud-break, whereas, for both sites, it is significantly different at veraison. Comparing all sites according to designed-systems, functional richness is significantly different between designed-systems at the two vine vegetative periods.

For all Designed-systems in a same site, bacterial abundance (Fig. 1B) is never significantly different between. At bud-break, bacterial abundance is between 28 and 45 ng/g of dry soil. For sites-systems Chaten_AB and Inger_AB, bacterial abundance is statistically different than the others sites-systems (respectively between 496-527 and between 65-141 ng/g of dry soil during veraison vine). One reason may be the number of years of organic farming in Inger_AB and Chaten_AB (more than 10 years) against less than 10 years for Ribeau_AB, and the others PI_Designed_system. Comparing all sites according to designed-systems, bacterial abundance is significantly different between designed systems only for veraison period.

Nitrogen mineralization kinetic (Nmin-kinetic) does not allowed to separate either PI and AB designed-systems, neither sites-systems (Fig. 1C).The top Nmin-kinetic is for Ribeau_PI site-system whereas the slowest one are for Ribeau_AB and Rouff_PI sites-systems. Rouff_Piopti, Inger_AB and Chante_AB sites-systems have touchily the same

Nmin kinetic.

It possible to separate PI and AB designed-systems according to PC2 axis of the PCA (Fig.1D): AB-systems under zero and PI-systems up to zero. Ribeauvillé-site is just on the y-axis for both systems AB and PI, i.e. there is no such difference between the two designed systems on this site. Inger_AB and Chante_AB are mainly explained by variables functional richness at bud-break and at veraison, whereas Rouff and Ribeau systems are mainly explained by bacterial abundance.




Fig. 1. Multicriteria results for all the sites-systems in Alsace, for 2014 vintage (France).

4 Conclusions

In this paper, we show that some of the analyzed BCN indicators are interesting to assess new designed-systems in different vineyard.

Acknowledgements. The authors thank the funders of these research programs: the ECOPHYTO and CASDAR programs of the French Department of Agriculture and Alsace region.

References

- Barlow, K., Bond, W., Holzapfel, B., Smith, J., and Hutton, R. (2009). Nitrogen concentrations in soil solution and surface run-off on irrigated vineyards in Australia. *Australian Journal of Grape and Wine Research* 15, 131-143.
- Guilpart, N., Metay, A., and Gary, C. (2014). Grapevine bud fertility and number of berries per bunch are determined by water and nitrogen stress around flowering in the previous year. *European Journal of Agronomy* 54, 9-20.
- Ingels, C. A., Scow, K. M., Whisson, D. A., and Drenovsky, R. E. (2005). Effects of cover crops on grapevines, yield, juice composition, soil microbial ecology, and gopher activity. *American Journal of Enology and Viticulture* 56, 19-29.
- Raath, P. J., and Saayman, D. (1995). Nitrogen mineralisation in vineyard soils of the Western Cape as affected by soil management practices. *South African Journal of Enology and Viticulture* 16, 7-13.
- Thiebeau, P., Herre, C., Doledec, A. F., Perraud, A., Panigai, L., Mary, B., and Nicolardot, B. (2005). Effect of soil covering on nitrogen availability of vineyards soils in champagne area. *Journal International Des Sciences De La Vigne Et Du Vin* 39, 163-177.

Proceedings of the 5th International Symposium for farming Systems Design

FSD5
Montpellier, September 7 - 10, 2015

Editors: Gritti Emmanuel S. – Wery Jacques
Cover design: Olivier Piau – Lisbeth Michel
Final edition: Gritti Emmanuel S.

Special thanks to the scientific committee's members for their commitment in the reviewing and editing processes