Resource management planning and ecological intensification to address climate challenges in French forestry

Jean-Francois Dhote, Barry Gardiner, Jean-Michel J.-M. Leban, Myriam Legay, Brigitte Musch, Christine Deleuze, Claudine Richter

▶ **To cite this version:**

Jean-Francois Dhote, Barry Gardiner, Jean-Michel J.-M. Leban, Myriam Legay, Brigitte Musch, et al.. Resource management planning and ecological intensification to address climate challenges in French forestry. Conférence franco-chinoise sur l'agro-écologie dans le contexte du changement climatique, Jun 2015, Pékin, China. pp.39 slides. hal-02794532

HAL Id: hal-02794532
https://hal.inrae.fr/hal-02794532

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Resource management planning and ecological intensification to address climate challenges in French forestry

Jean-François Dhôte, Barry Gardiner, Jean-Michel Leban (INRA), Myriam Legay, Brigitte Musch, Christine Deleuze, Claudine Richter (ONF)

Conférence franco-chinoise sur l'agro-écologie dans le contexte du changement climatique

Pékin, 3-5 juin 2015

3 Jun 2015

Objectives of the talk

- General background :
  - increasing world population → demand for wood-based products
  - forests provide an increasing range of products and services
  - forests are under pressure from other land-uses
  - supplying world's demand is a huge challenge to the sector
  - integrated response to climate change : adaptation, mitigation, ecosystem service regulation
- Climate change and the case of French temperate forests :
  - **natural regeneration**, large-size timber, **long rotations**
  - **imitation of nature** (Lorentz & Parade, 1837) → « *close-to-nature forestry* »
  - ➡ outline some challenges for forest **management planning**
  - ➡ define **ecological intensification** to meet these challenges

Highly diverse forest types

Oaks, Scots Pine (lowlands)

Beech

Fir, spruce, larch (mountains)



Maritime Pine



Mediterranean forests



Source : Memento (IGN, 2015)

1

Mitigate climate change : timber marketing, sawmilling efficiency, innovation for harvesting

Various technologies to value wood : from high-tech properties to energy

as building material ;
toll-station, motorway A89 (Limoges)

as bioenergy ;
Planoise district heating - 13 000 t/year (Besançon)

Favour wood design in emblematic buildings



Glue-laminated structure :
Simonin Frères
(Montlebon, France)


Adapt to sawmilling efficiency optimized
for Ø30-50cm logs
Siat sawmill (Urmatt, Alsace) :
700 000 m³/yr
©J.F. Dhôte, 2008
26/02/08 11:05

Conifer value curve reflects sawmilling process

Timber price (relative units)



Log diameter (cm)

Fir-Spruce - Vosges mean (sept. 2009-sept. 2010)

Source : ONF (Gamblin, 21/3/2011)

Max value

90% of Max value

*Opportunity for **shorter rotations** :*

- *reduce abiotic risks*
- *supply closer to industrial demand*

Innovation for harvesting

Sources : FCBA (E. Cacot, 2014) & ONF (E. Ulrich, 2014)

- prevent soil compaction : machine design and harvest optimal scheduling
- hardwood harvesting, work on slopes
- using data provided by harvesters

Avoir de nouveaux systèmes
techniquement et économiquement viables !

Integrated supply chain : better use of information linked to production processes



_2

Multi-scale, large-magnitude risks : prevent and mitigate damages

Local-scale impacts

Scattered Beech dieback at Fontainebleau (sept. 2011)

©J.F. Dhôte, 2011

Forest-scale impacts

©J.F. Dhôte, 2010

Regional-scale impacts → forestry-wood chain destabilization (eg storms)



Tempête Martin (1999) perte de 24 millions de m^3 de pin maritime



Tempête Klaus (2009) : perte de 43 millions de m^3 de pin maritime



Sources : Meredieu et al. (2014), Gardiner et al. (2010)



From regional to continental-scale disturbances



Mountain Pine Beetle outbreak in Western North America

Source : http://ngm.nationalgeographic.com/2015/04/pine-beetles/epidemic-map


Shortening rotation ages
to reduce abiotic risk sensitivity :
- storm damages
- drought-induced weakening
©S. Dhôte, 2011
how to conserve biodiversity of
old/senescent forest-cycle stages ?

Reducing water uptake by lower LAI : heavier, frequent thinnings



but… Disturbance & soil compaction
Which species install in the understory ?
Is this effective ?

©J.F. Dhôte, 2014

Favouring mixed stands…

©S. Dhôte, 2012

mixtures are eventually more productive, and more resistant against biotic hazards

3

Uncertainty and adaptation : explore/ combine ≠ diversification options

Uncertainty : for which climates to adapt ?



Anomaly of **summer temperature** 2100, models Aladin et LMDz scenario A1B

Source : M. Legay (2012)

Anomaly of **summer rainfall** 2100, models Aladin et LMDz scenario A1B

Changing species/genetic resources… : moving provenances polewards



- **(in northern hemisphere)**
  - vulnerability of populations at southern edge of distribution area
  - transfer seeds
  - plant in northern locations
- **Possible applications**
  - for **conserving genetic resources**
  - for **supporting local adaptation of existing species**

Changing species/genetic resources…



Changing production systems : e.g. product-targeted, short-rotation silviculture

Nombre de placettes par modalité

Species :	Fertilisation : Ash	Ash	Ca + Mg	No	No
	Soil preparation: Yes	No	Yes	Yes	No
Douglas fir	1 100 t/ha 3				
Douglas fir	1 600 t/ha 3	3	3	3	3
Douglas fir	2 000 t/ha 3				
Norway spruce	2 200 t/ha 3				
Cupressocyparis	1 600 t/ha 3				
Willow	2 000 t/ha 3	3	3	3	3
Black locust	2 000 t/ha 3				
Spontaneous vegetation	3	3		3	3

Source : ONF (Richter, 2014)

Planning forests with ≠ ways to regenerate

Special case : Oak forests

- Natural regeneration « *as usual* »
  - Special case : genetic resource conservation
- Same with very short rotation
- Planting southern provenances
- Planting related species to favour hybrids
- Introducing acclimated species
- Introducing exotic species

Planning forests with ≠ rotation ages

- Long rotations for *mainstream* forestry & conservation :
  - Standard rotation age : ≈ optimal silviculture
  - Longer rotation : ageing areas (delayed harvesting)
  - No rotation age : senescence areas and biological reserves (no more harvesting)
- Short rotations for specific management objectives :
  - product-oriented silvicultures to decrease harvesting pressure on *mainstream* forestry : bioenergy, small sawlogs…
  - increase adaptability : fasten genetic turn-over
  - handle species with present vitality ≠ long-term viability prognosis

4

Conclusions

Conclusions (1/2)

- Climate challenges **drive** forest management change :
  - Monitor/analyse local damages, anticipate large-scale crises
  - Increasingly rely on existing and new planted forests
  - Enhance resilience at ≠ levels of management systems
  - Achieve higher degree of forestry-wood chain integration
  - Optimize the use of the world's wood fibres and energy
  - Much more targeted & efficient use of forest land
- Some general rules for change ?
  - No obvious, no *one-size-fits-all* solutions : **diversify options**
  - **Multi-scale**, adaptive resource **management planning**
  - **Ecological intensification** (≠ close-to-nature forestry)

Conclusions (2/2)

- Multi-scale, adaptive resource management planning :
  - Combine ≠ ways of **diversification** : plantation/natural regeneration, mixtures, production systems, rotation ages…
  - Support **resilience** : shorter rotations, liming, low-impact harvesting, logistics (fight against storms, wildfires)…
  - **Information** : inventories, monitoring, product processing
  - Funding, priority setting and **investment** allocation
- Ecological intensification :
  - make **more efficient use** of natural and man-induced **cycles**…
  - of matter, energy and information…
  - in **ecosystems** as well as in the emerging **bioeconomy**

谢谢你的关注
Thank you for your attention
Merci

6

Annexes

French forests in a nutshell

- 16,5 millions hectares (30% of country area)
- + 8 M ha in French Guyana
- Expanding : 9 M ha in 1830, +87 000 ha/an (1980-2011)
- **74% private** owners, 17% communities, **9% State**
- 88% natural regeneration, **12% plantation**
- > 130 tree species
- Stock : 2,6 billion m^3, **64% hardwoods**, 36% conifers
- Summary of flux (average 2005-13)
  - Raw production : 91,5 Mm^3/yr
  - Mortality : 8,5 Mm^3/yr
  - **Harvest** : 41,4 Mm^3/yr (**50%** of net increment)
  - Stock increase : 41,4 Mm^3/yr

Source : IGN (2015)

Final harvest in highest-quality Oak Forests
rotation age : 180 to 250 years

Source : ONF (Jarret, 2014)

Natural regeneration

Source : ONF
(Jarret, 2014)
age : 1 yr

Young stand tending (mixture & competition control)

age : 25 yrs

Present best products : barrel manufacturing

Source : ONF
(Jarret, 2014)

Today, this is a very profitable production system,
Question :
very long rotations + exposition to climate/social risks :
how to diversify and increase resilience ?


Beech : 10% of growing stock
Rotation age : 120-180 yrs
North-east
North-west
©J.F. Dhôte, 2010

Silver fir : 8% of growing stock
Rotation age : 100-150 yrs
Uneven-aged management :
irregular shelterwood system (« *Plenterwald* »)
©J.F. Dhôte, 2010, 2014

Production systems based on plantation…

Maritime pine : 5% of growing stock

Rotation age : 40-50 yrs

Other conifers : 5%



Douglas fir : 4%

Rotation age : 50-80 yrs

Norway spruce : 8% of growing stock

Rotation age : 70-120 yrs

© C. Dhôte, 2005

©J.F. Dhôte, 2009

©J.F. Dhôte, 2015

Rotation age : 15-30 yrs

Hybrid poplar plantations :
1,2% of growing stock (30 M m
1,2% of area (180 000 ha)