

Folate content of cultivated and wild traditional leafy vegetables found in Nigeria

Shirley Isibhakhomen Ejoh, Faustina Dufie Wireko-Manu, David Page,

Catherine M.G.C. Renard

► To cite this version:

Shirley Isibhakhomen Ejoh, Faustina Dufie Wireko-Manu, David Page, Catherine M.G.C. Renard. Folate content of cultivated and wild traditional leafy vegetables found in Nigeria. Experimental Biology 2016, Apr 2016, San Diego, United States. 2016. hal-02794952

HAL Id: hal-02794952 https://hal.inrae.fr/hal-02794952

Submitted on 5 Jun2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés. **Folate content of cultivated and wild traditional leafy vegetables found in Nigeria AUTHORS:** Shirley Isibhakhomen **EJOH**^{1, 2, 3}, Faustina **DUFIE WIREKO-MANU**^{1, 2, 4}, David **PAGE**^{1, 2}, Catherine M.G.C. **RENARD**^{1,2}

1. INRA, UMR408 Sécurité et Qualité des Produits d'Origine Végétale, F-84000 Avignon, France

2. Avignon Université, UMR408 Sécurité et Qualité des Produits d'Origine Végétale, F-84000 Avignon, France.

3. Department of Human Nutrition, Faculty of Public Health, College of Medicine, University of Ibadan, Nigeria.

4. Department of Food Science and Technology, Kwame Nkrumah University of Science and Technology, Kumasi, Ghana.

Folate is an important micronutrient in the prevention of neural tube defects and cardiovascular disease. Given its importance, it is necessary to establish accurately the folate content foods that could contribute significantly to dietary intake. The aim of the study was to quantify the folate content in raw and cooked traditional leafy and non-leafy vegetables because there is a dearth of information on the folate content leafy vegetables found in Nigeria. Therefore the folate content of seven traditional leafy vegetables and one none leafy vegetable found in Nigeria were determined. The vegetables included: *Corchorus olitorius, Adansonia digitata, Abelmoscus esculentus (fruit), Abelmoscus esculentus (leaves), Amaranthus hybridus, Launaea taraxacifolia, Solanum macrocarpon, and Crassocephalum crepidioides.* They were further cooked in boiling water, until assessed as cooked according to Nigerian eating habits..

Folate is found in foods mainly as polyglutamate. The analysis of folate in the raw and cooked vegetable samples followed the process of extraction, deconjugation (using chicken pancreas deconjugase), derivatisation (a series of chemical reactions to convert all folate present in the deconjugated sample extracts, to 5-methyltetrahydrofolic acid (THF-5CH₃) monosodium glutamate and / or diglutamate), purification by affinity chromatography with Folate Binding Protein and quantification by HPLC equipped with fluorimetric detection. Folate content in the raw samples ranged from 183.4 µg/100g (FW) in *Corchorus olitorius* to 21.5 µg/100g (FW) in *Solanum macrocarpon* leaves; and in the boiled samples from 48.6 µg/100g (FW) in *Launaea taraxacifolia* to 8.5 µg/100g (FW) in *Solanum macrocarpon*. Loss of folate in the boiled vegetables varied from 46.6% in *Launaea taraxacifolia* to 88.4% in *Adansonia digitata*.

Traditional green leafy vegetables are good sources of folate. However cooking of the vegetables caused a considerable decrease in the folate content of the vegetables. Considering that most leafy vegetables are consumed in their cooked form in Nigeria, preparation methods of traditional leafy vegetables that would allow for optimal retention of folate are necessary.