

# Assessment of factors affecting genomic selection in populations of tomato

Coralie Picard, Janejira Duangjit, Mathilde M. Causse, Christopher Sauvage

### ▶ To cite this version:

Coralie Picard, Janejira Duangjit, Mathilde M. Causse, Christopher Sauvage. Assessment of factors affecting genomic selection in populations of tomato. 12. Solanaceae Conference, Oct 2015, Bordeaux, France. hal-02795275

HAL Id: hal-02795275 https://hal.inrae.fr/hal-02795275

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## The 12th Solanaceae Conference

October 25 -29, 2015 ENSEIRB Building, Talence

### Bordeaux, France



#### Assessment of factors affecting genomic selection in populations of tomato

Coralie Picard a, Janejira Duangjit a,b, Mathilde Causse a and Christopher Sauvage a

Marker-assisted selection is traditionally used to improve traits in crops. However, it requests very long cycles selection and still unable to capture loci of minor effect. Genomic selection (GS), a new tool for selecting elite plants in a breeding program by predicting the performance of traits of interest in application of statistical model, is an exciting alternative approach. GS has the potential to catch the effect of all markers underlying the genetic architecture of the studied traits.

In this context, we aimed at evaluating the use of GS into tomato to evaluate the prediction accuracy for 25 traits related to the fruit quality for three different type of population (RIL, MAGIC and GWA panel). We conducted a cross validation approach while estimating the relative weight of parameters (training and testing populations sizes, genetic composition, predictive statistical models, markers density, heritability of the trait as well as the type of population) onto the prediction accuracy.

Our results demonstrated that GS seems very powerful at predicting phenotypes values with accuracies ranging from 0.12 to 0.84 in the panel. Precisely, we conclude that (1) optimizing the training set the better the predictions are (+13% on average) (2) the larger the heritability of the trait value is, the better the predictions are (3) the statistical prediction models perform very similarly and (4) the optimal markers density is depending of the population LD.

Thus, applying GS in tomato seems very promising and could be used tomato breeding programs and produce high-quality varieties.

<sup>&</sup>lt;sup>a</sup> Institut National de la Recherche Agronomique, UR1052 GAFL, Génétique et Amélioration des Fruits et Légumes, 67 allée des chênes, CS60094, 84143 Montfavet cedex, France

<sup>&</sup>lt;sup>b</sup> Departments of Horticulture, Faculty of Agriculture, Kasetsart University, Bangkok, 10900 Thailand