



HAL
open science

Effects of a cultivar mixture on scab control in apple orchards

Laurent Brun, Arnaud Lemarquand, Gilles Orain, Christophe Gros, Freddy Combe, Frédérique Didelot, Claude Eric Parveaud, Christelle Gomez, Luciana Parisi

► **To cite this version:**

Laurent Brun, Arnaud Lemarquand, Gilles Orain, Christophe Gros, Freddy Combe, et al.. Effects of a cultivar mixture on scab control in apple orchards. 16. International conference on organic fruit growing, Feb 2014, Hohenheim, Germany. 17 p. hal-02795483

HAL Id: hal-02795483

<https://hal.inrae.fr/hal-02795483>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Effects of a cultivar mixture on scab control in apple orchard

L. Brun¹, A. Lemarquand², G. Orain², C. Gros¹, F. Combe¹,
F. Didelot³, C-E. Parveaud⁴, C. Gomez⁴, L. Parisi⁵

¹INRA-UERI, Domaine de Gotheron, Saint-Marcel-lès-Valence, France

²INRA - UE Horticole, Beaucouzé, France


³INRA - IRHS, Beaucouzé, France

⁴GRAB, Domaine de Gotheron, Saint-Marcel-lès-Valence, France

⁵INRA - UR Pathologie Végétale, France



Institut National de la Recherche Agronomique



17-19 February 2014



Why mixing apple cultivars on the same plot ?

- To reduce epidemics of aerial pathogens at the plot scale
- Mixing apple cultivars having different level of scab susceptibility can reduce scab epidemic (*e.g.* Bousset *et al.*, 1997)

- Hypothesis
 - a resource dilution effect
 - a physical barrier



resistant cv. or
low-susceptible

What kind of cultivar mixture ?

- Rvi6 gene overcome in some EU countries
 - If the resistance is overcome, mixing resistant and high susceptible cultivars could increase pathogen diversity (Trapman, 2006)
- Mixture of resistant + low susceptible cv

Aim of the study

Assess the efficiency of a **cultivar mixture**
within the row between
a **low-susceptible** and a **resistant** cultivar
on scab development
in **two** sites


efficiency scab development

assess

low-susceptible
cultivar mixture
resistant cultivar


Experimental orchard

- 2 experimental orchards
 - organic, SE France, Rhone Valley
 - IPM, W France, Loire Valley
- Planted in 2004



Experimental orchard

- 2 cultivars
 - Melrouge: low susceptibility to scab
 - Pitchounette: resistant to scab (*Rvi6*)
- Melrouge was planted in pure stands and in a within row mixture with Pitchounette



- Treatments were replicated 3 times according to a block design

Scab control and assessment

- No scab symptom or heterogeneous scab pressure until 2007
- No fungicide application against scab in 2008 and 2009
 - to permit the disease development
 - to make comparison easier
- Rvi6 gene was overcome in June 2008 at La Rétuzière
- Scab assessment
 - scoring on leaves in May and June
 - 80 shoots x 3 repetition x 2 treatments x 2 sites*
 - Scoring on fruits at picking
 - 20 fruits / tree on each tree*

Scab infection risks




Scab infection risks

		Primary inoculum	Angers	Light	Moderate	Severe	Total
Gotheron	2008	Low	1	11	2	6	20
	2009	High	0	4	3	0	7
La Rétuzière	2008	Low	4	7	7	5	23
	2009	High	1	5	5	7	18


Gotheron 2008 : beginning of scab development in the orchard

Scab shoot severity







- Low level of disease at Gotheron in 2008

- Effect of the cultivar mixture on number of scab lesions except in June 2009




Scab shoot severity



- Low level of disease at Gotheron in 2008
- Effect of the cultivar mixture on number of scab lesions except in June 2009

Incidence of scab at picking




	Percentage of scabbed fruits at picking	
	Gotheron - 2008	
	Melrouge	Pitchounette
Pure stand	14.9	-
Cultivar mixture	9.3	0.0
<i>P-value</i>	<i>0.0666</i>	

	Percentage of scabbed fruit	
	Gotheron - 2009	
	Melrouge	Pitchounette
Pure stand	82.2	-
Cultivar mixture	76.2	0.0
<i>P-value</i>	<i>0.0458</i>	

- Significant effect of cultivar mixture on fruit damages on Melrouge in 2008 and at Gotheron in 2009
- No significant effect of mixture at La Rétuzière in 2009 when the level of the disease is high and when Pitchounette is overcome

Incidence of scab at picking



	Percentage of scabbed fruits at picking			
	Gotheron - 2008		La Rétuzière - 2008	
	Melrouge	Pitchounette	Melrouge	Pitchounette
Pure stand	14.9	-	73.0	-
Cultivar mixture	9.3	0.0	60.5	0.1
<i>P-value</i>	0.0666		0.0663	

	Percentage of scabbed fruits at picking			
	Gotheron - 2009		La Rétuzière - 2009	
	Melrouge	Pitchounette	Melrouge	Pitchounette
Pure stand	82.2	-	98.3	-
Cultivar mixture	76.2	0.0	96.0	9.2
<i>P-value</i>	0.0458		0.4769	

- Significant effect of cultivar mixture on fruit damages on Melrouge in 2008 and at Gotheron in 2009
- No significant effect of mixture at La Rétuzière in 2009 when the level of the disease is high and when Pitchounette is overcome

Conclusions

- The overcoming of the resistance depends on:
 - quantity of inoculum
 - climatic conditions
 - cultural practices (spraying program, sanitation, cv mixture)
 - susceptibility of the cultivar
 - ➔ difficult to assess the relative importance of the cv mixture on the overcoming of the resistance
- Significant effect of the cv mixture observed:
 - between a low susceptible and a resistant cultivar
 - when the resistance is not overcome and the scab pressure is not too high
 - ➔ Extrapolation to commercial orchards tricky
 - ➔ cv mixture must be associated with other cultural practices

Thank you for your attention !

efficiency

scab development

assess

low-susceptible

cultivar mixture

resistant cultivar

Effects of a cultivar mixture on scab control in apple orchard

L. Brun¹, A. Lemarquand², G. Orain², C. Gros¹, F. Combe¹,
F. Didelot³, C-E. Parveaud⁴, C. Gomez⁴, L. Parisi⁵

¹INRA-UERI, Domaine de Gotheron, Saint-Marcel-lès-Valence, France

²INRA - UE Horticole, Beaucouzé, France


³INRA - IRHS, Beaucouzé, France

⁴GRAB, Domaine de Gotheron, Saint-Marcel-lès-Valence, France

⁵INRA - UR Pathologie Végétale, France



Institut National de la Recherche Agronomique



17-19 February 2014



Melrouge and Pitchounette mixture within the row



Melrouge fruit, 25 August 2011

