HAL open science

Control of N2O emissions by soil hydraulic functioning at the local scale

Catherine Hénault, Eva Rabot, Marine Lacoste, Isabelle I. Cousin

To cite this version:

Catherine Hénault, Eva Rabot, Marine Lacoste, Isabelle I. Cousin. Control of N2O emissions by soil hydraulic functioning at the local scale. Jahrestagung der Deutschen Bodenkundlichen gesellschaft, Sep 2015, Munchen, Germany. hal-02796090

HAL Id: hal-02796090

https://hal.inrae.fr/hal-02796090v1

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Control of N_2O emissions by soil hydraulic functioning at the local scale

Hénault C., Rabot E., Lacoste M., Cousin I.

2015/09/08
Deutsche Bodenkundliche Gesellschaft

INTRODUCTION

- N_2O emissions are largely controled by soil WFPS

- WFPS, combines informations on soil water content and on soil porosity. It is a proxy of the aerobic status of soil



Example of th *Fw* function in N_2O models (from Hénault and Germon, 2000)

- The relation between soil WFPS and N_2O emissions (denitrification) is exponential, N_2O emissions are very sensitive to little changes in soil WFPS

- This creates large uncertainties in N_2O assessments by simulations

INTRODUCTION

AIMS

- To investigate the « soil – water » interactions at the local scale and their consequences on soil N_2O emissions

CONTENTS

- Dynamics of N_2O emissions during wetting-drying cycles
- Soil porosity and N_2O emissions during wetting-drying cycles
- Modeling of new knowledges …

_01

Dynamics of N_2O emissions during wetting-drying cycles

Materials and Methods

Study site

- Cultivated with rape (*Brassica napus* L.)
- Classified as Glossic Retisol (WRB, 2014)
  - Silt loam texture

    clay = 13.7%, silt = 82.0%, sand = 4.3%
  - Soil organic carbon = 9.5 g kg^{-1}
- Bulk density = 1.43 g cm^{-3}

Soil sampling

Undisturbed soil core collected in a PVC cylinder (13.2-cm diameter by 7-cm height) from the surface horizon

Materials and Methods

N_2O measurements

- **Closed-chamber method**
- **N_2O monitoring: infrared correlation spectroscopy (N_2O Analyzer model 46i, Thermo Scientific), detection limit 0.02 ppm, concentration value recorded every minute**
- **Flux calculated linearly from the observed change in N_2O concentration during the first 10 min after the chamber was closed**



Materials and Methods

Hydric control

Multistep outflow system



Materials and Methods

Hydric control

- **Creation of two wetting-drying cycles with different rates of drying**

Multistep outflow system



Results and Discussion



Rabot et al. 2014 - SSSAJ

- Main peaks of N_2O emissions are observed
  - → not during the wetting phase
  - → but during the drying one
- The shapes of the peaks depend of the rate of the drying phase
- Peaks are attributed to
  - → N_2O dissolution during the wetting phase
  - → N_2O transport when soil pores are reconnected

⇨ To continue to investigate the effect of soil structure (porosity)

⇨ To introduce a transport module and a dissolution module into N_2O models

_02

Study of the soil porosity and N_2O emissions during wetting-drying cycles

Materials and Methods

X-ray computed tomography

Medical X-ray scanner: Siemens Somatom Definition AS

Acquisition time: 70 s

Voxel size: 316 × 316 × 100 µm

- **1 scan at the end of the wetting phase**
- **7 to 9 scans during the drying phase**

Alternating with N_2O measurements

Materials and Methods

Image analysis

- **Pre-processing:** ImageJ *(Rasband, 1997-2014)* and Avizo software







- Description of the pore network geometry : Image J software, C/C++ Quantlm library (Vogel, 2008)
  - Volume of air-filled and water-filled macropores
  - Euler number : connectivity of the air-filled pore space (vogel et al., 2010)
- Functionning description
  - Relative gas diffusion coefficient (DS/D0) from the segmented air-filled pore space

Results and discussion



Rabot et al. In Press – Vadoze Zone Journal

Results and discussion









Rabot et al. In Press – Vadoze Zone Journal

Results and discussion







Water saturated pores

Rabot et al. In Press – Vadoze Zone Journal

Results and discussion







Rabot et al. In press – Vadoze Zone Journal

_03

Modeling of this new knowledges

Materials and Methods

The NOE ALGORITHM

(Hénault et al., 2005 – Global Change Biology)

$$N_2O = (N_2O)_{denit} + (N_2O)_{nit}$$

- $denit = K \cdot F_T \cdot F_{NO3-} \cdot F_{WFPS}$

  (Hénault and Germon, 2000)

- $nit = f \cdot (H_m \cdot [NH_4^+] \cdot T)$

Biological parameters

Environmental functions and parameters







Conceptual proposition (NOE_{GTE})

A transport module

- 1D diffusion equation of Fick
- Use of the Millington and Quirk model (1961)

A dissolution module

- A gas-liquid equilibrium according to Henry's law



Fig. 1 Simulated nitrous oxide flux with time for water-filled pore space in the range [0, 1]. At time = 0 h, initial conditions were applied. For the next time increments, the nitrous oxide production module, and the gas transport and equilibrium modules were activated

Rabot et al. 2015 – Biogeochemistry

Experimental evaluation

- Measurements and simulation of N_2O emissions in laboratory on undisturbed soil cores submitted to wetting/drying cycles with different rates



Fig. 5 Measured (*black squares*) and 95 % confidence interval of simulated N_2O fluxes (*grey areas*). Sample names refer to the number of decreasing pressure heads applied during one wetting–drying cycle (Fast: 3 values, Medium: 4 values, Slow: 6 values, *BD* bulk density in g cm^{-3}). *Yellow areas* represent the drying phases, and *white areas* the wetting phases. (Color figure online)

Rabot et al. 2015 – Biogeochemistry

CONCLUSIONS

- Great interest of dynamic studies : they have revealed peaks of N_2O emissions during the drying phase of wetting-drying cycles

- To continue to characterise soil structure, pore soil network. This is probably a key for understanding soil emissions dynamics (our study could be improved by higher resolution tomography) and intensity.

- To continue to define some physical indicators of soil structure and to introduce them in modeling