



HAL
open science

Plant interactions as biotic drivers of intraspecific variability in leaf litter traits and decomposability of a foundation tree species (*Quercus petraea*)

Ludovic Henneron, Matthieu Chauvat, Frédéric Archaux, Marthe Akpa-Vinceslas, Fabrice Bureau, Yann Dumas, Laurent Mignot, Francois Ningre, Sandrine Perret, Claudine Richter, et al.

► To cite this version:

Ludovic Henneron, Matthieu Chauvat, Frédéric Archaux, Marthe Akpa-Vinceslas, Fabrice Bureau, et al.. Plant interactions as biotic drivers of intraspecific variability in leaf litter traits and decomposability of a foundation tree species (*Quercus petraea*). Colloque international de la Société française d'écologie SFECOLOGIE 2016, Oct 2016, Marseille, France. 2016. hal-02796562

HAL Id: hal-02796562

<https://hal.inrae.fr/hal-02796562>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Plant interactions as biotic drivers of intraspecific variability in leaf litter traits and decomposability of a foundation tree species (*Quercus petraea*)

Ludovic Henneron^{1,2}, Matthieu Chauvat¹, Frédéric Archaux³, Marthe Akpa-Vinceslas¹, Fabrice Bureau¹, Yann Dumas³, Laurent Mignot¹, François Ningre⁴, Sandrine Perret³, Claudine Richter⁵, Philippe Balandier³ and Michaël Aubert¹


¹ Normandie Univ, UNIROUEN, IRSTEA, ECODIV, FR-76000 Rouen, France; ² UREP, INRA, Clermont-Ferrand, Auvergne, France; ³ IRSTEA, Unit Research on Forest Ecosystems, Domaine des Barres, F-45290 Nogent-sur-Vernisson, France; ⁴ INRA-LERFoB, 54280 Champenoux; ⁵ ONF, Research & Development department, Boulevard de Constance, 77300 Fontainebleau, France
ludovic_henneron@hotmail.com



Context & scientific issues

- **Leaf litter traits and decomposability** are increasingly recognized as an 'extended' phenotype affecting plant community dynamics and ecological processes through plant-litter-soil feedbacks.
- **Intraspecific trait variability** can be an important source of functional diversity. While the role of genetic variability and environmental conditions, *i.e.* mostly soil abiotic factors, as sources of intraspecific variation in leaf litter traits and decomposability has been explored, very few studies have focused on the effect of **biotic factors such as plant neighbor identity and abundance and plant-plant interactions**.
- **Removal experiments** can be useful tools to manipulate species abundance in real ecosystems and to study the effect of altering plant interactions. Hence, the **removal of oak canopy trees** can affect oak leaf litter quality by altering: competition for light among trees (H_1); competition for nutrients among trees (H_2); competition for nutrients between trees and coexisting understory plants (H_3), either related to abundance changes (H_{3a}) or shifts in species composition (H_{3b}) of understory plants; soil abiotic properties alterations (H_4).

Aim of the study: to assess how plant interactions control intraspecific variation in oak leaf litter traits and decomposability



Material & methods

Study sites & treatments:



- **Experimental manipulation of oak abundance** through the removal of oak canopy trees using silvicultural-thinning regimes of contrasting intensity (Fig. 2)
- 5 Forest sites located throughout northern France on **contrasted pedoclimatic contexts** (Fig. 3):
 - Oceanic to continental temperate climate
 - Very acidic (moder) to slightly acidic (mull) soil conditions
- Utilization of two complementary **experimental networks**:
 - The Scientific Interest Group - French national cooperative for data on forest growth - oak group (GIS-Coop)
 - The oak network of the Forest and Wood Resource Laboratory (LERFoB)
- 19 sessile oak (*Quercus petraea*) stands studied covering a wide gradient of:
 - **Oak abundance** → 3 to 44 m²·ha⁻¹ (oak canopy tree basal area used as a proxy)
 - **Forest successional stage** → 18 to 171 years old stands
 - ✓ **Aggradation** stage (< 90-year-old) → intense tree height growth and plant competition
 - ✓ **Biostatic** stage (>90-year-old) → maturity of trees




Oak leaf litter sampling and analyses:

- Sampling by **litter traps** (Fig. 4)
- A large set of 26 **litter chemical and biochemical traits** to characterize oak leaf litter quality
- Decomposability of oak leaf litter determined using **standardized *ex situ* decomposition bioassays** (Fig. 5)

Results & discussion



- Oak leaf litter quality strongly declined with tree removal in early forest successional stage, which had important negative consequences for litter decomposability
 - Litter became poorer in nutrients such as N and Mg and richer in secondary metabolites such as total phenolics, condensed tannins and lignin (Fig 6, 7a)
 - Importantly, litter N loss switched from N release to N immobilization (Fig 7b)
- Variance partitioning indicated that oak abundance explained as much variation in oak leaf litter traits as oak age and twice as much as soil abiotic factors
 - This underlines that biotic factors can be more important sources of intraspecific variation in tree leaf litter traits than more commonly recognized soil abiotic factors



- The decline of oak leaf litter quality induced by tree removal was most likely driven by a shift in understory plant community composition (H_{3b} , Fig. 8).
 - This could be due to an increased competition with a more abundant and faster-growing understory plant community with an higher leaf N content
 - We propose that this plasticity of leaf litter traits could be an adaptive strategy of ectomycorrhizal tree species consisting in 'short-circuiting' the N cycling through plant-litter-soil feedbacks in order to preserve their pre-empted nutrient pool of the capture by competing understory plants in the context of treefall gaps.

Conclusions & broader implications

- Our large-scale, multi-site study involving long-term oak canopy tree removal provides clear experimental evidence that **biotic factors such as plant-plant interactions can be strong drivers of plasticity in leaf litter traits**, with important consequences for litter decomposability.
- This finding contributes to the emerging view that **phenotypic plasticity** is fundamentally related to biotic interactions for sessile organisms, especially for long-lived and large plant species such as trees.
 - Taking this source of functional diversity into account could help us to better understand plant community dynamics and ecological processes in terrestrial ecosystems.

Acknowledgements:
This study was funded by a grant provided by the Upper-Normandie region via the Wide Area Network Research (TERA: Territory, Environment, Risk, Agronomy) and by the French Ministry of Ecology, Sustainable Development and Energy (Convention No. 10-VBGE-BGF-3-CVS-081, BGF program, IMPREBIO Research Project). We greatly thank P. Delporte, M. Le Gall, C. Abgrail, J. Lainé, A. Follin, M. Colombel and S. Mascré for laboratory, technical and field support. We also thank M. Makkonen for sharing their data, J. Messier for their advice in statistical analysis as well as N. Fanin for their helpful comments.
We also thank the 'French data cooperative on forest growth' (GIS Coop) and the 'oak network of the Forest and Wood Resource Laboratory' (LERFoB), supported by AgroparisTech, INRA, ONF and Irstea with the financial support of the French Ministry of Agriculture and Forest, for the management of several sites studied and for providing data.