



HAL
open science

Intercropping of lentil and spring wheat inscreases productivity in organic farming

Loic Viguiet, Laurent Bedoussac, Etienne-Pascal Journet, Eric Justes

► To cite this version:

Loic Viguiet, Laurent Bedoussac, Etienne-Pascal Journet, Eric Justes. Intercropping of lentil and spring wheat inscreases productivity in organic farming. 14. European Society of Agronomy Conference 2016-ESA14, Sep 2016, Edinburg, United Kingdom. , 2016. hal-02796589

HAL Id: hal-02796589

<https://hal.inrae.fr/hal-02796589>

Submitted on 20 Jun 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Context

- Interest in **Lentil** (*Lens culinaris*) is **steadily growing** for human consumption in France & EU
- Demand in organic lentil is higher than the supply in France
- Organic lentil **productivity is low and instable**
- Growing lentil in mixture could be a way to **support farmers** in fulfilling consumers demand



Objectives

1. Analyse the **functioning** of lentil-spring wheat mixture or intercrops (IC)
2. Evaluate their performances for **yield improvement/stabilization** and **income**
3. Find the **best combinations of cultivars and sowing densities** for organic farmers





Materials & Methods

- Field experiment was conducted in 2015 under **organic agricultural conditions**
- 4 cultivars of lentils (L) and 2 of spring wheat (W) were grown as sole crops (SC)
- **32 bi-specific intercrops** (4 lentil cv. x 2 wheat cv. x 4 seeding ratios) were evaluated
- Both species were **sown** and **harvested** together at the lentil ripeness stage
- Grain yield, yield components and Nitrogen acquired were measured
- Grain sorting of bi-specific mixture was carried out in post-harvest



Results



- **Total intercrop yield ≈ wheat sole crop > lentils sole crop**
- **Wheat always dominated** lentil particularly for the 67%L/33%W and the 100%L/50%W density ratios
- Lentil grain yield in IC lower than in SC due to strong **interspecific interactions** produced by wheat on lentil
- **In SC, Anicia, Rosana and Flora** yields were higher than Beluga
- **In IC, Flora and Rosana** yields were higher than Anicia and Beluga
- Cv. Beluga yield was **similar in IC than in SC**
- **The higher the wheat density, the lower the lentil grain yield** without increasing the total IC yield

Conclusions & perspectives

- Lentil-spring wheat intercrops are of great interest in organic farming to **improve yields & income** (incl. grain sorting)
- **Intercrop efficiency** due to the **complementarities of both species for the use of abiotic resources**
- To favour the lentil yield (the most profitable crop), **wheat density** must remain low (< 33%) and lentil sown at 100%
- **Harvest efficiency** of lentils in IC compared to SC seems better, but other results are needed to verify the agronomic and economic advantages obtained in this experiment

