



**HAL**  
open science

## Regulation of tomato ascorbate levels: a candidate gene involved in fruit nutritional value and stress tolerance

Noé Gest, Vincent Truffault, Hicham El Airaj, Cecile Garchery, Gisèle Riqueau, Mathilde M. Causse, Hélène Gautier, A. Jimenez, Rebecca Stevens

### ► To cite this version:

Noé Gest, Vincent Truffault, Hicham El Airaj, Cecile Garchery, Gisèle Riqueau, et al.. Regulation of tomato ascorbate levels: a candidate gene involved in fruit nutritional value and stress tolerance. XVIIIth Eucarpia Meeting, Vegetable section, Tomato Working group, Apr 2014, Avignon, France. 1 p., 2014. hal-02796653

**HAL Id: hal-02796653**

**<https://hal.inrae.fr/hal-02796653v1>**

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



## P007 – Rebecca Stevens

### ***Regulation of tomato ascorbate levels: a candidate gene involved in fruit nutritional value and stress tolerance***

Gest, N. (1), Truffaut, V. (1,2), El Airaj, H. (1), Garchery, C. (1), Riqueau, G. (1), Causse, M. (1), Gautier, H. (2), Jimenez, A. (3), Stevens, R. (1)

(1) INRA, Génétique et Amélioration des Fruits et Légumes, Montfavet, France

(2) INRA, Plantes et Systèmes de Culture Horticole, Avignon, France

(3) CEBAS-CSIC Stress Biology and Plant Pathology, Murcia, Spain

Ascorbate is a powerful antioxidant in plants, and its levels are an important quality criterion in commercial species. Factors influencing these levels include environmental variations, particularly light, and the genetic control of its biosynthesis, recycling and degradation. Our previous work has identified QTL for fruit vitamin C content and a candidate gene, encoding a monodehydroascorbate reductase (MDHAR) isoform, an enzyme catalysing reduction of the oxidized radical of ascorbate, monodehydroascorbate, to ascorbate. The activity of this enzyme is also correlated with fruit tolerance to post-harvest chilling stress in introgression lines containing the vitamin C QTL. In order to understand the role of this gene in fruit physiology and metabolism we produced transgenic lines in different genetic backgrounds. We show that the isoform encodes a cytosolic and peroxisomal MDHAR, and that this enzyme negatively regulates ascorbate levels in certain genotypes of *Solanum lycopersicum* (tomato). Transgenic lines overexpressing MDHAR show a decrease in ascorbate levels in leaves, whereas lines where MDHAR is silenced show an increase in these levels in both fruit and leaves. Furthermore, the intensity of these differences is light-dependent. The unexpected effect of this MDHAR on ascorbate levels cannot be explained by changes in the expression of Smirnoff-Wheeler pathway genes, or the activity of enzymes involved in degradation (ascorbate peroxidase) or recycling of ascorbate (dehydroascorbate reductase and glutathione reductase) suggesting a previously unidentified mechanism regulating ascorbate levels. The impact of under-expression of this gene in processing tomato on fruit post-harvest physiology has also been examined.

**Keywords:** ascorbate, post-harvest, QTL, redox