



## A SNP map of the European sea bass genome based on a meiotic gynogenetic family

Münnever Oral, Julie Colléter, Michaël Bekaert, John B. Taggart, Christos Palaiokostas, Brendan J. Mcandrew, Marc Vandeputte, Béatrice Chatain, David J. Penman

### ► To cite this version:

Münnever Oral, Julie Colléter, Michaël Bekaert, John B. Taggart, Christos Palaiokostas, et al.. A SNP map of the European sea bass genome based on a meiotic gynogenetic family. Aquaculture Europe 2014, Oct 2014, San Sebastian, Spain. hal-02797976

**HAL Id: hal-02797976**

**<https://hal.inrae.fr/hal-02797976>**

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**A SNP map of the European sea bass  
(*Dicentrarchus labrax*) genome based on a  
meiotic gynogenetic family**

**Aquaculture Europe 2014 - San Sebastián**  
**16<sup>th</sup> October 2014**

**Münevver Oral**, Julie Colléter, Michaël Bekaert, John B Taggart, Christos Palaiokostas, Brendan J McAndrew, Marc Vandeputte, Béatrice Chatain, Heiner Kuhl, Richard Reinhardt and David J Penman

**Institute of Aquaculture, Stirling, UK**  
**AQUAEXCEL Workpackage 9: Isogenic Fish Lines**



## I. Background

- Importance of Clonal Lines
- Production of such lines
- Why Meiotic Gynogenesis
- Aim of the present study

## II. Material & Methods

- Production of Meiotic Gynogenetics
- ddRAD lib construction
- Genetic Linkage Map (R/OneMAP) & Physical Map

## III. Results & Discussion

- ddRAD reads summary
- Genetic Linkage Map
- Physical Map
- Crossover hotspots


## IV. Conclusion & Suggestions

### **Clonal Lines = Genetically Identical Individuals**

- Homogeneity (powerful for the investigation of complex genetic traits, decreases the variation in experiments)
- Standardisation to the research –refined experimental design (Aquaculture related research point of view)
- Speed of generation (2 subsequent production cycles via Gynogenesis-G1 or Androgenesis-A1)
- Eliminates deleterious recessive alleles, reveals genetic variation for many traits
- QTL (Quantitative Trait Loci) identification and whole genome sequencing projects

# I. Background: Production of Clonal Lines

## MITOTIC GYNOGENESIS




**Gynogenesis (G):** All maternal inheritance  
**Androgenesis (A):** All paternal inheritance

Spontaneous meiotic gynogenetics can be a problem when producing isogenic clonal lines via mitotic gynogenesis.

FAO, *Inbreeding and Broodstock Management*  
 Chapter 6, Chromosome Set Manipulations.

# I. Background: Meiotic Gynogenetics




## Aim of the study

was to define **telomeric markers** by constructing a SNP based genetic linkage map in *D. labrax* by facilitating the power of high-throughput sequencing to differentiate between meiotic to mitotic gynogenetics.



Centromeric Loci will  
be homozygous



Intermediate markers  
will show varying level  
of heterozygosity

VERY Telomeric markers  
will be heterozygous



These can distinguish  
between meiotic and  
mitotic gynogenetics



## Next Generation Sequencing (NGS)

Drowned in next generation sequencing data



[www.biocomicals.com](http://www.biocomicals.com), Alper Uzun, PhD



## II. Material & Methods: Production of Meiogynogenetics



Ifremer Experimental Aquaculture Station (Palavas-les-Flots, France).

UV irradiation device:

Four lamps above  
Four lamps below (12 W, 254 nm,  
Vilber-Lourmat, Marne-la-Vallée,  
France)

Sperm irradiation (1:20, v/v in SGSS) for 8 minutes was applied for total dose of  $326 \text{ mJ/cm}^2$  which corresponds to the dose of  $32000 \text{ erg/mm}^2$  described by (Peruzzi & Chatain, 2000).

## II. Material & Methods

### DNA extraction

- One Meiotic Gynogenetic *D. labrax* family:
  - Sire (♂) and dam (♀): Fin Clips
  - 80 offspring: Whole larvae
- REALpure kit (REAL Laboratories Spain)
- Nanodrop Spectrophotometry
- Agarose Gel Electrophoresis
- Final DNA concentration assessment: Quantica qPCR thermal cycler by using dsDNA fluorescent dye.

### Double digest-ddRAD library preparation and sequencing

OPEN ACCESS Freely available online




#### Double Digest RADseq: An Inexpensive Method for *De Novo* SNP Discovery and Genotyping in Model and Non-Model Species

Brant K. Peterson\*, Jesse N. Weber, Emily H. Kay, Heidi S. Fisher, Hopi E. Hoekstra

Department of Organismic & Evolutionary Biology, Department of Molecular & Cellular Biology, Museum of Comparative Zoology, Harvard University, Cambridge, Massachusetts, United States of America

- Peterson et al., 2012


## II. Material Methods: Double Digest RAD seq (ddRAD)



Sequence & Analyse



### III.Results: Sequencing & RAD tag summary



From purified gDNA to the genotypes takes 1 week.

EBI Sequence Read Archive (SRA) with the accession number of: ERP006697.

**Raw Reads:** All reads have been produced by sequencer.

**Filtered reads:** Reads with right barcodes & adapters combination. Stacks package (Catchen et al., 2011).



## II. Material Method: Genetic Linkage Map

- 340 male specific markers (no paternal inheritance detected)
- **R/OneMap** v2.0-4 (Margarido *et al.*, 2007) & **TMAP** v1.1 (Cartwright *et al.*, 2007).
  - 804 female heterogametic Input file: Outcross (Kosambi function)
  - *rf.2pts*
  - *LOD=4, max.rf=0.5 (draft seabass genome assembly)*
  - *order.seq (ser,rcd,rec,ug)*
  - *safe & force*
  - *rf.graph.table*
- *Final genetic map: Genetic-Mapper v0.7 (Bekaert, 2012).*

Table 2: Notation used to identify markers and genotypes

	crosstype	Parent			Offspring		
		Cross		Observed bands	Observed bands	Segregation	
A		1	<i>ab</i> × <i>cd</i>	<i>ab</i> × <i>cd</i>	<i>ac, ad, bc, bd</i>	1:1:1:1	
		2	<i>ab</i> × <i>ac</i>	<i>ab</i> × <i>ac</i>	<i>a, ac, ba, bc</i>	1:1:1:1	
		3	<i>ab</i> × <i>co</i>	<i>ab</i> × <i>c</i>	<i>ac, a, bc, b</i>	1:1:1:1	
		4	<i>ao</i> × <i>bo</i>	<i>a</i> × <i>b</i>	<i>ab, a, b, o</i>	1:1:1:1	
B	B <sub>1</sub>	5	<i>ab</i> × <i>ao</i>	<i>ab</i> × <i>a</i>	<i>ab, 2a, b</i>	1:2:1	
	B <sub>2</sub>	6	<i>ao</i> × <i>ab</i>	<i>a</i> × <i>ab</i>	<i>ab, 2a, b</i>	1:2:1	
	B <sub>3</sub>	7	<i>ab</i> × <i>ab</i>	<i>ab</i> × <i>ab</i>	<i>a, 2ab, b</i>	1:2:1	
C		8	<i>ao</i> × <i>ao</i>	<i>a</i> × <i>a</i>	<i>3a, o</i>	3:1	
D	D <sub>1</sub>	9	<i>ab</i> × <i>cc</i>	<i>ab</i> × <i>c</i>	<i>ac, bc</i>	1:1	
		10	<i>ab</i> × <i>aa</i>	<i>ab</i> × <i>a</i>	<i>a, ab</i>	1:1	
		11	<i>ab</i> × <i>oo</i>	<i>ab</i> × <i>o</i>	<i>a, b</i>	1:1	
		12	<i>bo</i> × <i>aa</i>	<i>b</i> × <i>a</i>	<i>ab, a</i>	1:1	
		13	<i>ao</i> × <i>oo</i>	<i>a</i> × <i>o</i>	<i>a, o</i>	1:1	
	D <sub>2</sub>	14	<i>cc</i> × <i>ab</i>	<i>c</i> × <i>ab</i>	<i>ac, bc</i>	1:1	
		15	<i>aa</i> × <i>ab</i>	<i>a</i> × <i>ab</i>	<i>a, ab</i>	1:1	
		16	<i>oo</i> × <i>ab</i>	<i>o</i> × <i>ab</i>	<i>a, b</i>	1:1	
		17	<i>aa</i> × <i>bo</i>	<i>a</i> × <i>b</i>	<i>ab, a</i>	1:1	
		18	<i>oo</i> × <i>ao</i>	<i>o</i> × <i>a</i>	<i>a, o</i>	1:1	

Refer to: Wu *et al.* (2002) Simultaneous maximum likelihood estimation of linkage and linkage phases in outcrossing species.

## Physical Map


- 804 female heterogametic SNP markers as well as 11 microsatellites have been assigned in to European sea bass draft genome contigs across 24 LGs.
- Has been used to define recombination points.

MB\_microsatellites position on physical map  
Marker\_ID,Physical\_Map\_(Genome\_Location),Physical\_Map\_Position\_(bp)

Dla0003,LG19,8127283  
Dla0006,LG6,25269003  
Dla0104,LG2,24689183  
Dla0105,LG8,10127559  
Dla0106,LG2,25784997  
Dla0112,LG5,12681454  
Dla0119,LG14,8349079  
Labrax17,LGX,16504896  
Labrax29,LG18-21,9732294  
Labrax3,LG13,6001586  
Labrax8 ,LG16,2733963

(Chistiakov *et al.*, 2005)  
(García De León *et al.*, 1995)

### III.Results: Meiogynogenetic Linkage Map (R/OneMap)




Basic karyotype,  $2n=48$

LGs	Number of Markers	Length (cM)
LG1A	45	78.04
LG1B	29	51.30
LG2	30	61.83
LG3	24	22.79
LG4	34	44.10
LG5	38	68.19
LG6	26	55.59
LG7	26	72.31
LG8	31	47.92
LG9	23	46.00
LG10	30	34.68
LG11	42	54.26
LG12	29	47.25
LG13	27	54.03
LG14	31	66.67
LG15	37	61.31
LG16	37	49.89
LG17	45	55.03
LG18-21	15	30.23
LG19	30	56.96
LG20	46	45.82
LG22-25	39	62.70
LG24	19	39.07
LGX	31	45.05
<b>Total</b>	<b>764</b>	<b>1252.02</b>

### III.Results: Final *D.labrax* Meiotic Genetic Map

Empty circles: Homozygote  
Black dots: Heterozygote







# LG11

Empty circles: Homozygote  
Black dots: Heterozygote

Overall

Individual progeny



Average  $0.90 \pm 0.08$  crossover  
per chromosome arm.


## IV. Conclusion

- 764 SNP markers have been identified and mapped from the single meiogynogenetic *D. labrax* family with 79 offspring+parents.
- Crossover points per chromosome arm have been identified per linkage group with an average of  $0.90 \pm 0.08$  crossover per chromosome arm.
- Particularly those markers closer to telomere are of interest to differentiate between meiotic and mitotic gynogenesis for the reliable production of isogenic clonal lines (meiotic gynogenetics need to be detected and eliminated).
- Cost effective and quick (<£10 per individual, up to 100 individuals and >500 SNPs in one sequencing lane)
- Genotyping by (RAD) sequencing rather than isolating and assaying these individually or in multiplexes.

## V. Future work

- Application to development of Isogenic clonal lines in:
  - Seabass (*Dicentrarchus labrax*),
  - Common carp (*Cyprinus carpio*),
  - the Atlantic salmon (*Salmo salar*)
  - (Nile Tilapia (*Oreochromis niloticus*))
- Gametic inactivation in each species using UV irradiation and perhaps X-rays (androgenesis – chromosome fragments?).
- Distinguishing between meiotic and mitotic gynogenetics
- Genetic Linkage map.
- Verification of isogenic lines through RADseq (starting from outbred founders (parents of G1/A1 fish) to G1/A1 clone founders (with biparental control) and to isogenic lines.

# Experimental Approaches by using Genotyping by Synthesis



Baird *et al.*, 2008 (RAD seq) & Peterson *et al.*, 2012 (ddRAD seq)





Would you like to find out more?



**AQUAEXCEL INDUSTRY WORKSHOP:**  
Research Infrastructures: adding value to  
European aquaculture industry

Friday, 17<sup>th</sup> October  
Kicking off at 10.30am  
Room 11 (Exhibition Area)

**See you there!**





# Contact us

## Thank you for your attention

Email: [d.j.penman@stir.ac.uk](mailto:d.j.penman@stir.ac.uk)

Website: [www.aquaexcel.eu](http://www.aquaexcel.eu)

Tel: +44 (0)1786 467901



# DISCLAIMER



The research leading to these results has received funding from the European Union's Seventh Framework Programme (FP7/2007-2013) under grant agreement no 262336. This publication reflects the views only of the author, and the European Union cannot be held responsible for any use which may be made of the information contained therein.