



HAL
open science

Towards functional genomics of transcription factor genes associated to growth and wood formation in maritime pine

Jean-François Trontin, Concepción Avila, Sandrine Debille, Fernando de La Torre, Jorge El-Azaz, Belen Pascual, Francis Canlet, Caroline Teyssier, Nathalie Boizot, Claire C. Le Mette, et al.

► To cite this version:

Jean-François Trontin, Concepción Avila, Sandrine Debille, Fernando de La Torre, Jorge El-Azaz, et al.. Towards functional genomics of transcription factor genes associated to growth and wood formation in maritime pine. ProCoGen final open conference, Promoting Conifer Genomic Resources, Nov 2015, Orléans, France. , pp.1, 2015. hal-02798385

HAL Id: hal-02798385

<https://hal.inrae.fr/hal-02798385>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



ProCoGen final open conference

30th November – 2nd December 2015

Promoting Conifer Genomic Resources

- *Structural genomics*
- *Functional genomics*
- *Comparative genomics*
- *Translational genomics*

ProCoGen Training and dissemination workshops

3rd – 4th December 2015

- *Practicalities of marker and genome-assisted selection*
- *From our labs to your forests*
- *Transfer of genomic tools to breeding programs*

Centre de Conférences d'Orléans, France

Towards functional genomics of transcription factor genes associated to growth and wood formation in maritime pine

Jean-François Trontin¹, Concepción Avila², Sandrine Debille¹, Fernando de la Torre², Jorge El-Azaz², Belén Pascual², Francis Canlet¹, Caroline Teyssier³, Boizot N³, Le Metté C³, Lesage-Descauses M-C³, Denilson da Silva Perez⁴, Rafael Cañas², Grégoire Le Provost⁵, Christophe Plomion⁵, Luc Harvengt¹, Philippe Label⁶, Marie-Anne Lelu-Walter³, Francisco Cánovas²

¹FCBA, Pôle Biotechnologie et Sylviculture Avancée, 71 route d'Arcachon, Pierroton, 33610 Cestas, France

²Universidad de Málaga, Dept. Biología Molecular y Bioquímica, Facultad de Ciencias e Instituto Andaluz de Biología, Campus de Teatinos S/N, 29071 Málaga, Spain

³INRA, UR 0588 Unité Amélioration, Génétique et Physiologie Forestières, 2163 avenue de la Pomme de Pin, CS 4001, Ardon, 45075 Orléans Cedex 2, France

⁴FCBA, Pôle IntechFibres, Domaine Universitaire, CS 90251, 38044 Grenoble Cedex, France

⁵INRA-Université de Bordeaux, UMR 1202 BIOGECO, 69 route d'Arcachon, Pierroton, 33612 Cestas, France

⁶INRA-Université Blaise Pascal, UMR A547 PIAF, Les Cèzeaux, 24 Avenue des Landais, 63177 Aubière cedex, France

Maritime pine is a major forest tree in southern Europe for sustainable delivery of bio-based products through advanced plantation forestry of improved varieties. Early selection is needed to reduce breeding cycle and accelerate variety deployment in the context of climate change. Both gene(s) and genome-wide information should provide opportunities for predictive, marker-assisted selection. Reverse genetics, defined as ectopic expression or silencing of candidate genes can be useful for functional dissection of traits of interest. Functional genomics of 9 transcription factor (TF) genes associated to growth and wood formation (*MYB1,8,14,20,23*, *DOF5*, *MADBOX4*, *NACx*, *NACataf*) was initiated in maritime pine through genetic transformation of embryogenic tissue. Twelve TF constructs designed for constitutive overexpression (OE) or silencing (RNAi) were studied: 6 constructs (batch 1) from previous projects (*MYB1*-RNAi, *MYB8*-OE/RNAi, *MYB14*-RNAi, *DOF5*-OE/RNAi) as well as 3 constructs for putative gene targets of *MYB* (*CAD*-RNAi) and *DOF5* (*GS1a*-OE, *GS2*-OE); and 6 constructs (batch 2) obtained during ProCoGen (*MYB23*-OE, *MYB20*-OE, *MADBOX4*-OE, *NACx*-OE/RNAi, *NACataf*-OE).

Phosphinothricin-resistant lines could be cryopreserved for all but one (*MYB20*-OE) constructs. Transformation rate was estimated in the range 9.6-22.0 (OE) or 0.4-18.0 (RNAi) transgenic lines per gram embryogenic tissue (batch 1). Somatic embryos were obtained from 1-3 lines per construct (25 lines in total) and successfully converted to plants (batch 1). Acclimatization rates were similar to controls and transgenic plants were confirmed for most lines. Putative adverse effect of *MYB14*-RNAi, *DOF5*-RNAi, and *MYB8*-OE on transformation rate was observed. *DOF5*-RNAi apparently stimulated germination rate. Plant growth data and morphology are available for constructs from batch 1 after up to 12 (*MYB1*-RNAi, *MYB8*-OE/RNAi, *DOF5*-OE/RNAi, *GS2*-OE, *GS1a*-OE) or 42 months (*MYB14*-RNAi, *CAD*-RNAi) growth. Both transgene copy number and targeted gene expression data are available from transgenic plants obtained from *MYB14*-RNAi and *CAD*-RNAi lines. Transgenics and controls from batch 1 were sampled at age 16 (*MYB1*-RNAi, *MYB8*-OE/RNAi, *DOF5*-OE/RNAi) or 70 months (*MYB14*-RNAi, *CAD*-RNAi). Molecular characterization of *MYB1,8,14* and *DOF5* plant material has been performed by qPCR and transcriptomic analysis using microarray is underway. Wood analyses have been initiated for *MYB14*- and *CAD*-RNAi plants (in progress).

This work received funding from the European Community's Seventh Framework Programme (FP7/2007-2013, Grant Agreement N°289841-PROCOGEN).