



# Advances in multiple pests modelling: IPSIM (Injury Profile SIMulator) a hierarchical qualitative modelling framework to predict injury profiles as a function of cropping practices, soil, climate and field environment.

Marie-Hélène Robin, Jean-Noel Aubertot

## ► To cite this version:

Marie-Hélène Robin, Jean-Noel Aubertot. Advances in multiple pests modelling: IPSIM (Injury Profile SIMulator) a hierarchical qualitative modelling framework to predict injury profiles as a function of cropping practices, soil, climate and field environment.. PURE Congress, Jan 2015, Poznan, Poland. 2015. hal-02798964

**HAL Id: hal-02798964**

**<https://hal.inrae.fr/hal-02798964>**

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.





# INJURY PROFILE SIMULATOR (IPSIM), A MODELLING PLATFORM TO DESIGN QUALITATIVE MODELS PREDICTING INJURY PROFILES AS A FUNCTION OF CROPPING PRACTICES AND PRODUCTION SITUATIONS



ROBINMH<sup>1,2</sup>, AUBERTOT JN<sup>2</sup>

1 INPT, INP-EIPurpan, 75 voie du Toec BP57611, 31076Toulouse cedex 3, France  
2 INRA, AGIR, UMR 1248, 31326 Castanet-Tolosan, France  
E-mail: mh.robin@purpan.fr



## INTRODUCTION


In order to reduce the reliance of cropping systems on pesticides, tools to help "vertical integration" (combination of several control methods) and "horizontal integration" (simultaneous management of several pests) in Integrated Pest Management strategies are needed. Wheat is the main arable crop in France in terms of cultivated area. We propose **an innovative modelling framework in order to help design qualitative models to represent the impact of cropping practices, soil, climate and field environment on injury profiles caused by multiple pests** (plant pathogens, weeds and animal pests). The basic principles of the approach and an application to wheat, the main arable crop in Europe in terms of cultivated area is presented here.

## MATERIALS AND METHODS


IPSIM is a simple generic hierarchical qualitative modeling platform based on the DEXi software (Bohanec, 2009). The main assumption is that each injury profile that can be observed in a given field only depends on the associated cropping system and part of the production situation (soil, climate and field environment). Figure 1 represents the conceptual scheme of IPSIM-WHEAT. In order to provide a proof of concept, the method was first applied to a single disease: eyespot on wheat (Robin et al., 2013). IPSIM-WHEAT-EYESPOT, a sub-model of IPSIM-WHEAT, was designed and its predictive quality was assessed (Figure 2). Utility functions were defined using available knowledge in the literature. It must be stressed that no calibration was performed. A large dataset (525 site-years) was used to assess the predictive quality of IPSIM-WHEAT-EYESPOT.



© Bayer CropScience




**Figure 1. Schematic representation of IPSIM-WHEAT.** In blue, green and orange: production situation components affecting the injury profile of a given wheat field. The injury profile is the output variable of IPSIM-WHEAT. Its input variables are embedded within the three following components: soil and climate, field's environment, cropping practices.



**Figure 2. Structure of IPSIM-WHEAT-EYESPOT.**


## RESULTS AND DISCUSSION



**Figure 3. Distribution of differences between observed and simulated final eyespot incidence (0-20%, 20-40%, 40-60%, 60-80%, 80-100%; 525 fields, over 9 years and 19 French regions)**

The predictive quality of IPSIM-WHEAT-EYESPOT is satisfactory, even if slightly biased. Half of the simulated values are similar to observations and 80.5% differ at most by one class (Figures 3, 4). In addition to this evaluation, the predictive quality of IPSIM-WHEAT-EYESPOT was characterised with statistical criterion using the 5 class centers as representants of simulated final incidence classes, and the observed final incidence values instead of classes. IPSIM-WHEAT-EYESPOT proved to fairly represent the effects of cropping practices as well as the regional and annual variability of the disease (Efficiency = 0.51, Root Mean Square Error of Prediction = 24%; bias = 5.0%). Other sub-models are under construction.

The platform IPSIM appears to be an innovative tool to design qualitative models predicting injury profiles to help design agroecosystems less reliant on pesticides



**Figure 4. Confusion matrix of the IPSIM-Wheat-Eyespot model and marginal distributions. Numbers in italic are overall percentages.**