



**HAL**  
open science

# Transcriptional and chromatin-based control of effector gene expression in the plant pathogenic fungus *Leptosphaeria maculans*

Jessica L. Soyer, Mennat El Ghalid, Jonathan Grandaubert, Marie-Helene Balesdent, Lanelle R Connolly, Michael Freitag, Thierry T. Rouxel, Isabelle Fudal-Grolier Fudal

## ► To cite this version:

Jessica L. Soyer, Mennat El Ghalid, Jonathan Grandaubert, Marie-Helene Balesdent, Lanelle R Connolly, et al.. Transcriptional and chromatin-based control of effector gene expression in the plant pathogenic fungus *Leptosphaeria maculans*. ECFG13. European conference on fungal genetics, Apr 2016, Paris, France. hal-02799514

**HAL Id: hal-02799514**

**<https://hal.inrae.fr/hal-02799514>**

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Transcriptional and chromatin-based control of effector gene expression in the plant pathogenic fungus *Leptosphaeria maculans*

Jessica L. Soyer<sup>1</sup>, Mennat El Ghalid<sup>1</sup>, Jonathan Grandaubert<sup>1</sup>, Marie-Hélène Balesdent<sup>1</sup>, Lanelle R. Connolly<sup>2</sup>, Michael Freitag<sup>2</sup>, Thierry Rouxel<sup>1</sup> and Isabelle Fudal<sup>1</sup>

<sup>1</sup>INRA / AgroParisTech, UMR 1290 BIOGER, Thiverval-Grignon, France.

<sup>2</sup>Department of Biochemistry and Biophysics, Oregon State University, Corvallis, USA.

### Abstract

Plant pathogens secrete an arsenal of small secreted proteins (SSPs) acting as effectors that modulate host immunity to facilitate infection. In fungi, SSP-encoding genes are often located in particular genomic environments and show waves of concerted expression during plant infection. To date, little is known about the regulation of their expression. *Leptosphaeria maculans*, a fungus responsible for stem canker of oilseed rape, has a bipartite genome structure alternating gene-rich and transposable element (TE)-rich regions. TE-rich regions, which encompass one third of the genome, are enriched in putative effector genes that present the same expression pattern (no or a low expression level during *in vitro* growth and a strong over-expression *in planta*). On these bases, we investigated the determinism of the concerted effector gene expression testing two hypotheses: (i) are one or several common regulators involved in the control of the concerted expression of effector genes? and / or (ii) are TE-rich regions targets of reversible chromatin modifications that affect the regulation of effector gene expression?

To identify putative regulators of effector gene expression, we established the repertoire of TFs of *L. maculans* and identified TFs only found in *L. maculans* genome or specifically induced during infection. We performed functional analyses on several TFs and showed that StuA plays a major role in infection and expression of effector genes in *L. maculans*. We also investigated the involvement of one histone modification, histone H3 lysine 9 tri-methylation (H3K9me3) in chromatin-based regulation of concerted effector gene expression. For this purpose, we silenced expression of two key players in heterochromatin assembly and maintenance, *HP1* and *Kmt1*, by RNAi. Whole genome oligoarrays performed on silenced-*HP1* and silenced-*Kmt1* transformants revealed an over-expression of SSP-encoding genes in TE-rich regions during *in vitro* growth. That increase of expression was associated with a reduction of H3K9 tri-methylation at two SSP-encoding gene loci. These data strongly suggest that a chromatin-based control, mediated by HP1 and KMT1, represses the expression of at least part of the effector genes located in TE-rich regions during growth in axenic culture. Our hypothesis is that changes of lifestyle and a switch toward pathogenesis lift chromatin-mediated repression, allowing a concerted expression of effector genes mediated by one / or several TF(s).