



**HAL**  
open science

## **GWAS of tomato response to water deficit: focus on major fruit quality traits**

Elise Albert, Vincent Segura, Justine Gricourt, Romain Novaretti, Yolande Carretero, Esther Pelpoir, Julien Bonnefoi, Laurent Derivot, Mathilde Causse

### ► **To cite this version:**

Elise Albert, Vincent Segura, Justine Gricourt, Romain Novaretti, Yolande Carretero, et al.. GWAS of tomato response to water deficit: focus on major fruit quality traits. 13. Solanaceae Conference, Sep 2016, Davis, United States. 2016. hal-02799533

**HAL Id: hal-02799533**

**<https://hal.inrae.fr/hal-02799533v1>**

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



**Albert E.**<sup>a</sup>, Segura V.<sup>b</sup>, Gricourt J.<sup>a</sup>, Novaretti R.<sup>a</sup>, Carretero Y.<sup>a</sup>, Pelpoir E.<sup>a</sup>, Bonnefoi J.<sup>c</sup>, Derivot L.<sup>c</sup>, Causse M.<sup>a</sup>  
<sup>a</sup>INRA, UR1052 GAFL, 84143 Montfavet, France | <sup>b</sup>INRA, UR0588 AGPF, F-45075 Orléans, France | <sup>c</sup>GAUTIER Semences, F-13630 Eyragues, France

## Context

- Water scarcity is a crucial constraint for agriculture productivity.
- **Water deficit** can improve **tomato flavour** by concentrating the **major taste compounds** in the fruit, but the right balance must be found to **limit yield loss**. [1]
- There is a **high genetic variability** for response to water deficit in cultivated tomato (*S. lycopersicum*), in particular in **small fruited accessions** (*S. l. cerasiforme*). [2]

## Objectives

Using small fruited accessions, we aimed to (1) characterize the pattern of **genotype by watering regime interactions** at the **phenotypic & genotypic levels**, and (2) identify the **major loci and genes** involved in tomato fruit quality variation under water deficit.

## Materials & Methods

- **141 highly diverse small fruited accessions**, among which **4 were fully re-sequenced** [5]
- **Greenhouse** experiment with **two watering conditions**
  - well watered: WW
  - water deficit: WD = 40% WW
- **11 fruit traits**: fruit number, fresh weight (FW), dry matter (DMW), soluble sugar (SSC), ascorbic acid (VitC), glucose, fructose, pH, malic & citric acid content
- **6,501 SNP** genotyped over the genome (SOLCAP array)
- **Genome Wide Association (GWA)**:
  - multi loci mixed model: **MLMM** [3] (highly polygenic traits)
  - multi trait mixed model: **MTMM** [4] (G x W tests)
- Publicly available **expression data** (tomato genome consortium)

## Results




Fig 1 | Variance partitioning

← The trait variations resulted from **large genotypic** effects and **medium genotype by watering regime interaction** effects.  
 (W = watering, Gr = genetic group, G = genotype)




Fig 2 | Fruit quality & Yield variations  
 $\Delta = (WD-WW)/WW$

↑ **Fifty accessions** (with small to medium fruit size) had **both improved fruit SSC and maintained fruit number** under WD.




Fig 3 | Marker-trait associations according to their type

↑ A total of **81 associations** were identified, **51% were specific** to one condition (**WD** or **WW**) and **18% were interactive** with the watering regime.




Fig 4 | Manhattan plot & LD heatmap

↑ An association was detected on chr11 for **Fructose under WD**, including in its linkage disequilibrium block **5 genes**, all **expressed in tomato fruit**. Among these genes, one encodes a 'Neutral Invertase' with a **non-synonymous variants** between the 4 re-sequenced accessions of the GWA collection.

## Take home messages

- ✓ Tomato **quality** could be **improved** under deficit irrigation **while maintaining yield**.
- ✓ The underlying genetic architecture relies on **numerous loci** with **small effects** and **varies with water availability**.
- ✓ **Genes** related to **sugar metabolism** were identified under some associations and could control the variations observed.

## Funding



## References

- [1] Ripoll et al. 2014      [4] Segura et al. 2012  
 [2] Albert et al. 2016    [5] Causse et al. 2013  
 [3] Korte et al. 2013

## Contact

[AlbertElise2@gmail.com](mailto:AlbertElise2@gmail.com)