



**HAL**  
open science

## Metabolic profiling of transgenic lines silenced for an MDHAR gene and impact on ascorbate redox state and carbon metabolism

Vincent Truffault, Guy Costagliola, H el ene Gautier, Rebecca Stevens

► **To cite this version:**

Vincent Truffault, Guy Costagliola, H el ene Gautier, Rebecca Stevens. Metabolic profiling of transgenic lines silenced for an MDHAR gene and impact on ascorbate redox state and carbon metabolism. 18. EUCARPIA Meeting of the Tomato Working Group, Apr 2014, Avignon, France. 1 p. hal-02799822

**HAL Id: hal-02799822**

**<https://hal.inrae.fr/hal-02799822>**

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.



## T009 - Vincent Truffault

### *Metabolic profiling of transgenic lines silenced for an MDHAR gene and impact on ascorbate redox state and carbon metabolism.*

Truffault, V. (1,2); Costagliola, G. (2); Gautier, H. (2); Stevens, R. (1)

(1) INRA, Génétique et Amélioration des Fruits et Légumes, Montfavet, France

(2) INRA, Plantes et Systèmes de Culture Horticole, Avignon, France

Ascorbate is a key molecule involved in numerous cellular processes in plants and is an essential antioxidant protecting the cell from reactive oxygen species (ROS) damage in case of environmental stress. The cellular concentration of ascorbate depends on its biosynthesis, recycling and degradation: these are under genetic control and closely related to environmental conditions.

The recycling pathway controls the redox state of the ascorbate pool and is especially important during stress responses. Recycling of oxidized ascorbate to the reduced form occurs via two enzymes monodehydroascorbate reductase (MDHAR) and dehydroascorbate reductase (DHAR). Transgenic lines silenced for an MDHAR gene in a cherry tomato cultivar have been generated. Our previous results have shown that the activity of MDHAR is correlated with reduced ascorbate and fruit firmness in tomatoes. The objective of this study was to determine whether limiting turnover of the ascorbate pool may affect central metabolism of tomato plantlets and fruits.

MDHAR modified plantlets showed a slower early growth and fruit size was also affected. Silencing MDHAR induced a modification of carbohydrate metabolism affecting particularly sugar concentrations. The ascorbate pool and ascorbate degradation products were also modified. Transgenic lines showed other strong variations of metabolites compared to wild type.

Ascorbate concentration was measured by spectrophotometry. Metabolite profiling of transgenic lines was provided by a GC-MS based survey of tomato metabolites. Results are discussed and shed light on the impact of a modified recycling pathway on metabolism in plantlets and fruits of tomato.

**Keywords:** ascorbate-recycling, metabolic profiling, redox state