



Island river levelling: a favorable situation to explore ground beetles (Coleoptera : Carabidae) ability to recolonize pioneer riparian habitats

Olivier Denux, Emmanuelle Dauffy-Richard, Jean-Pierre Rossi, Sylvie Augustin

► To cite this version:

Olivier Denux, Emmanuelle Dauffy-Richard, Jean-Pierre Rossi, Sylvie Augustin. Island river levelling: a favorable situation to explore ground beetles (Coleoptera : Carabidae) ability to recolonize pioneer riparian habitats. 10. European Congress of Entomology, Aug 2014, York, United Kingdom. 1p., 2014. hal-02800002

HAL Id: hal-02800002

<https://hal.inrae.fr/hal-02800002>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Island river levelling: a favorable situation to explore ground beetles (Coleoptera : Carabidae) ability to recolonize pioneer riparian habitats

Olivier Denux¹, Emmanuelle Dauffy-Richard(†)², Jean-Pierre Rossi³, Sylvie Augustin¹

1: INRA Orléans, UR0633 Zoologie Forestière, France 2: IRSTEA Nogent-sur-Vernisson, UR EFNO, France 3: INRA Montpellier, UMR CBGP, France


Introduction

The Loire, the last wild river in Western Europe, undergoes recurrent maintenance operations of bed to improve flow and to avoid reduction of the main stem. In autumn 2012, the territorial collectivities carried out levelling of one island in the Natural Reserve of Saint-Mesmin (Centre region, France). We took advantage of this work to launch a long term survey of the island recolonization by plants, birds, beavers and insects. We report here the impact on insect populations taking ground beetles (Coleoptera : Carabidae) as a model. We compared species (communities) before and just after the levelling.

Methods


Habitats

In 2012, before levelling, we identified 8 pioneer habitats on the island Loire valley. After levelling, 4 pioneer habitats were found and selected on the island.



Ground beetles sampling

We used pitfall traps to sample ground beetles between June and September 2012 – 2013 in the habitats under study.



Analyses

Species richness

We report the observed species richness and its bootstrap estimator with the associated standard error in each pioneer habitats (Colwell & Coddington, 1994).

Indicator species


We explored the indicator value of each species with regards to the habitats surveyed in 2012 by means of the IndVal index developed by Dufrêne & Legendre (1997).

We present here IndVal values >0.25 associated to probabilities <0.001.


Results

Species richness and abundance

We identified 36 species in 2012 (869 individuals) and 17 species in 2013 (1022 individuals). Only 15 species were found again after levelling. For each habitat, the specific richness decreased sharply after levelling.




The species absent in 2013 were occasionally encountered before levelling. The riparian species *Principidum punctulatum* and *Lionychus quadrimaculatus* were both dominant before and after levelling.



Indicator species

We identified 9 characteristic species of different habitats before levelling in 2012.



Conclusions

The pioneer island habitats have an important species richness. A decrease of the species richness was observed just after levelling, however this decline concerned infrequent species (1 to 5 individuals) and may be explained by a lower sample intensity in 2013 as compared to 2012. The predominant species in 2012 were also abundant after levelling works.

Three species, *Chaenellus vestitus*, *Cicindela hybrida* and *Principidum punctulatum*, typical of riparian communities on Loire river and characteristic of wet gravel, dry sand and mudflat were found before and just after levelling.