



HAL
open science

Unusual evolutionary mechanisms to escape Effector-Triggered-Immunity in the fungal phytopathogen *Leptosphaeria maculans*

Clémence Plissonneau, Bénédicte B. Ollivier, Isabelle Fudal-Grolier Fudal,
Thierry T. Rouxel, Marie-Helene Balesdent

► **To cite this version:**

Clémence Plissonneau, Bénédicte B. Ollivier, Isabelle Fudal-Grolier Fudal, Thierry T. Rouxel, Marie-Helene Balesdent. Unusual evolutionary mechanisms to escape Effector-Triggered-Immunity in the fungal phytopathogen *Leptosphaeria maculans*. ECFG13. European conference on fungal genetics, Apr 2016, PARIS-LA VILLETTE, France. p.490, 2016. hal-02800552

HAL Id: hal-02800552

<https://hal.inrae.fr/hal-02800552>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER SESSION ABSTRACTS
Session CS6 Ecological and population genomics
CS6W19

Wednesday 6th April
14:00 - 16:00

PLISSONNEAU Clémence (1), OLLIVIER Bénédicte (1), FUDAL Isabelle (1), ROUXEL Thierry (1), BALESDENT Marie-Hélène (1)
(1) -BIOGER INRA, AgroParisTech, Thiverval-Grignon, France

Unusual evolutionary mechanisms to escape Effector-Triggered-Immunity in the fungal phytopathogen *Leptosphaeria maculans*

Leptosphaeria maculans is responsible for the stem canker (phoma) disease of oilseed rape (*Brassica napus*). AvrLm3 and AvrLm4-7, two avirulence effector genes of *L. maculans*, are involved in an unusual relationship: the presence of AvrLm4-7 induces the Rlm7-mediated resistance but suppresses the Rlm3-mediated recognition (Plissonneau et al., 2015; doi: 10.1111/nph.13736). To investigate this relationship, we assessed the sequence diversity of AvrLm3 in *L. maculans* populations. The analysis of more than 200 isolates revealed a high level of allelic polymorphism for this gene, but no deletion event nor inactivating mutations were found. This observation contrasts with the presence/absence or RIP-(Repeat-induced-point-mutation) inactivation polymorphisms usually found for other *L. maculans* avirulence genes and questioned the role of AvrLm3 in fungal fitness. Two distinct mechanisms responsible for the «double virulent» phenotype (i.e. virulent toward both Rlm3 and Rlm7, «a3a7») were identified. In isolates displaying an inactivation of AvrLm4-7, amino acid changes in AvrLm3 were responsible for the virulent phenotype towards Rlm3. However, 56% of the a3a7 isolates displayed an avirulent allele of AvrLm3, combined with point mutations in AvrLm4-7. Such situations allow the fungus to escape Rlm7-mediated resistance while maintaining the suppression of the AvrLm3 phenotype effective. They also allow the fungus to escape to the recognition by two resistance genes while keeping the secretion of the two corresponding effector proteins. The complex evolutionary mechanisms displayed by *L. maculans* to escape Rlm3-mediated resistance while preserving AvrLm3 sequence integrity, along with the reduced virulence in isolates silenced for AvrLm3 confirmed the importance of this effector in pathogenicity towards *B. napus*.

This work was funded by the INRA SMaCH metaprogram «K-masstec» and the CTPS project «ICOSCOP»
