



HAL
open science

GWAS reveals candidate regions for metabolic traits in tomato, *Solanum lycopersicum*

Guillaume Bauchet, S. Grenier, N. Samson, J. Bonnet, L. Grivet, E. Bolygo, C. Baxter, Mathilde M. Causse

► **To cite this version:**

Guillaume Bauchet, S. Grenier, N. Samson, J. Bonnet, L. Grivet, et al.. GWAS reveals candidate regions for metabolic traits in tomato, *Solanum lycopersicum*. XVIIIth Eucarpia Meeting, Vegetable section, Tomato Working group, Apr 2014, Avignon, France. 1 p., 2014. hal-02800694

HAL Id: hal-02800694

<https://hal.inrae.fr/hal-02800694>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



T024 - Guillaume Bauchet

GWAS reveals candidate regions for metabolic traits in tomato, *Solanum lycopersicum*

Bauchet, G. (1,2), Grenier, S. (2), Samson, N. (2), Bonnet, J. (2), Grivet, L. (2), Bolygo, E. (3), Baxter, C. (3) Causse, M. (1)

(1) INRA GAFL, Montfavet, France

(2) Syngenta Seeds, Saint-Sauveur, France

(3) Syngenta Jealott's Hill International Research Centre, Bracknell, United Kingdom

Modern breeding can impact crop cultivars volatile fruit content, inducing low volatile profile and therefore low organoleptic quality ([Klee and Tieman 2013](#)). Genome-wide association studies have been successful in identifying genes involved in polygenic traits notably in tomato, and can be a useful way towards fruit quality improvement. Genotyping arrays ([Sim, Durstewitz et al. 2012](#); [Viquez-Zamora, Vosman et al. 2013](#)) are available and enable GWAS for traits of interest. We present an association panel including 70 wild relative accessions (*S. pimpinellifolium*), 170 admixed accessions (*S. lycopersiforme*) and 50 domesticated accessions (*S. lycopersicum*). GWAS analysis was conducted using single and multi-locus analysis (Yu et al., 2006; Segura et al., 2012) with 12000 SNP markers and a set of sugar-related, amino-acids as well as a broad range of volatiles. The present study is the first one in tomato reporting associations for a large set of volatiles at the genome scale. We found significant associations for 84 loci with a total of 17 traits including glucose, malate or phenylacetaldehyde. Identified loci were also concordant with previously published quantitative trait loci (guaiacol), while new loci were identified (phenylacetaldehyde). These results (1) provide a list of candidate loci to be functionally validated and (2) provide a powerful approach for finding genetic variants that can be directly used for fruit quality improvement and deciphering the genetic architecture of complex traits.

References

- Klee, H. J. and D. M. Tieman (2013). "Genetic challenges of flavor improvement in tomato." *Trends in genetics : TIG* 29(4): 257-262
- Segura, V., B. J. Vilhjalmsson, et al. (2012). "An efficient multi-locus mixed-model approach for genome-wide association studies in structured populations." *Nature Genetics*, 44(7): 825-830.
- Sim, S.-C., G. Durstewitz, et al. (2012). "Development of a Large SNP Genotyping Array and Generation of High-Density Genetic Maps in Tomato." *PLoS ONE*, 7(7): e40563.
- Viquez-Zamora, M., B. Vosman, et al. (2013). "Tomato breeding in the genomics era: insights from a SNP array." *BMC Genomics*, 14(1): 354.
- Yu, J., G. Pressoir, et al. (2006). "A unified mixed-model method for association mapping that accounts for multiple levels of relatedness." *Nature Genetics*, 38(2): 203-208.

Keywords: tomato, GWAS, metabolome, volatiles