



HAL
open science

Wheat effector assisted breeding for resistance to fungal pathogens

Marc-Henri M.-H. Lebrun, Thierry T. Langin, Thomas T. Kroj, J. Cockram, R. Oliver, G. Kema, R. Valade, S. Praud, V. Laurent, L. Duchalais, et al.

► To cite this version:

Marc-Henri M.-H. Lebrun, Thierry T. Langin, Thomas T. Kroj, J. Cockram, R. Oliver, et al.. Wheat effector assisted breeding for resistance to fungal pathogens. 9.International symposium on Septoria diseases of cereals, Apr 2016, Paris, France. p.123, 2016. hal-02800809

HAL Id: hal-02800809

<https://hal.inrae.fr/hal-02800809v1>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

POSTER SESSION ABSTRACTS
Session 5: Host genetics and resistance breeding
FRI-20

LEBRUN Marc Henri (1), WEAB CONSORTIUM
(1) BIOGER, INRA, AgroParisTech, Thiverval-Grignon, France

Wheat effector assisted breeding for resistance to fungal pathogens

WEAB consortium: Thierry LANGIN, INRA GDEC, Clermont-Ferrand, France; Thomas KROJ, INRA BGPI, Montpellier, France; James COCKRAM, NIAB, UK ; Richard OLIVER, CCDM, Australia; Gert KEMA, PRI, Netherlands; Romain VALADE, Arvalis, Thiverval-Grignon, France ; Sébastien PRAUD, BIOGEMMA, Clermont-Ferrand, France; Valérie LAURENT, Florimond-Desprez, Cappelle en Pévèle, France; Laure Duchalais, RAGT, Louville la Chenard, France; Volker LEIN, CETAC, Estrées-Saint-Denis, France

The discovery that fungal effector proteins are important for infection represents a novel opportunity for controlling plant diseases. Use of fungal effectors for resistance breeding is a game-changing technology creating opportunities and innovative methods to identify novel resistances to fungal diseases in plants. These methods are amenable to high throughput phenotyping. The recent availability of high-density genetic marker coverage of the wheat genome allows the mapping of novel resistances identified through such high throughput phenotyping. We are using necrotrophic protein effectors from *Parastagonospora nodorum* (Pn) and toxic proteins from *Fusarium graminearum* (Fg) and *Zymoseptoria tritici* (Zt) to detect resistance genes/QTLs in wheat. Complementary strategies will be used to detect a large array of resistance mechanisms to fungal effectors. Recombinant necrotrophic protein effectors and toxic proteins are produced in yeast and the purified proteins are delivered into wheat leaves by syringe infiltration. Symptom development is scored few days after infiltration. Screening of 220 elite French wheat cultivars with Pn ToxA, 1 and 3 has highlighted a large number of cultivars resistant to the 3 necrotrophic effectors, and only few cultivars sensitive to the three effectors, suggesting that previous breeding for field resistance to Pn (1960-1980) has led to the accumulation of insensitivity alleles. To validate this hypothesis, we are currently pathotyping these wheat cultivars with a French Pn isolate producing Tox1 and 3. Mapping of loci controlling insensitivity to Pn necrotrophic effectors and resistance to Pn isolate will be performed using genome-wide association analyses. This project will facilitate plant breeding efforts to select for resistance to important fungal pathogens by providing a toolkit of bio-molecular markers.