



HAL
open science

Study of expression heterogeneity of flagellar genes in an insect pathogenic bacterium *Xenorhabdus* using single cell technologies

Anne Lanois, Grégory Jubelin, Sylvie Pages, Alain Givaudan

► To cite this version:

Anne Lanois, Grégory Jubelin, Sylvie Pages, Alain Givaudan. Study of expression heterogeneity of flagellar genes in an insect pathogenic bacterium *Xenorhabdus* using single cell technologies. 5. International conference on analysis of microbial cells at the single cell level, Nov 2011, Carry-le-Rouët, France. 2011. <hal-02803492>

HAL Id: hal-02803492

<https://hal.inrae.fr/hal-02803492v1>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

Study of expression heterogeneity of flagellar genes in an insect pathogenic bacterium *Xenorhabdus* using single cell technologies




Anne Lanois, Grégory Jubelin, Sylvie Pagès and Alain Givaudan

Laboratoire DGIMI UMR INRA-UM2 1333 Université Montpellier 2, 34095 Montpellier cedex 05 (lanois@univ-montp2.fr)




Introduction. *Xenorhabdus nematophila* is a motile Gram-negative bacterium that forms a mutualistic association with a nematode and their association is highly pathogenic for a wide range of insect larvae. Upon insect entry, *Xenorhabdus* is released by its nematode partner, escapes insect defence responses, invades tissues and hemolymph (the insect equivalent of blood) and secretes virulence factors involved in the death of insect. Among the regulatory networks involved in the virulence of *Xenorhabdus*, the flagellar cascade (cf figure) controls the production of flagellum, an extracellular appendage required for bacterial motility, and exoenzymes such as protease, lipase and hemolysins¹.



X. nematophila : a motile and flagellated bacterium

In this study, we investigated flagellar gene expression dynamics at single-cell resolution using flow cytometry. Promoter sequence of *X. nematophila flgB* gene (class II gene) was first inserted into pPROBE-*gfp*[AAV], a highly stable vector to create a transcriptional fusion with the reporter gene *gfp*[AAV]². This *gfp* gene version encodes an unstable GFP with a half-life of ~60 min in *E. coli*. Destabilized GFP was used as an expression monitoring system since it allows studies of fine-tuned transcriptional process. This construction was then transferred into *X. nematophila* and *flgB* expression was analysed.




Références: (1) Lanois A, Jubelin G, Givaudan A (2008) Mol Microbiol 68: 516-533.
(2) G. Jubelin, S. Pagès, A. Lanois, M. H. Boyer, S. Gaudriault, J.-B. Ferdy and A. Givaudan (2011). Environmental Microbiology. May;13(5):1271-84.


Bimodal expression of P*flgB*-*gfp*[AAV] fusion in *X. nematophila*

Cell sorting
was performed with exponential grown culture using a FACSaria cell sorter during 2 hours in PBS buffer.

Flow cytometry analysis



Fluorescence microscopy




...each subpopulation is able to produce both states.

Expression heterogeneity remains temporal.

Sorted cells analysis



A/ qRT-PCR :

RNA purification from 10⁷ cells using Qiagen microprep RNeasy kit



B/ Pathology test :

200 bacteria of each population were injected in insect larvae hemolymph.



flgB⁻ sorted cells displayed an hyper-virulent phenotype in our insect model *Spodoptera*. In contrast, flgB⁺ sorted cells have a delayed virulence suggesting that the production of flagella is deleterious to the bacterial virulence.

- ➔ (i) This result confirms the differential level of *flgB* gene expression observed with the *gfp* reporter gene.
- (ii) The "OFF" population displays lower levels of class II and class III mRNA, but not class I.

Conclusion and perspectives

- This work reveals a bimodal expression of *flgB* flagellar gene in *X. nematophila*.
- Previous experiment using a class III gene (*fliC*) gave the same results. These data mean a mixed population of flagellated and non-flagellated bacteria occurs at the mid-exponential growth phase.
- Because heterogeneity (or phenotypic noise) is often associated with positive feedback loop in gene networks, we are currently examining the involvement of flagellar regulators in this phenomenon.