



High or low fructose? Consequences for sugar metabolism in peach fruit

Elsa Desnoues, Yves Y. Gibon, Valentina Baldazzi, Veronique Signoret, Michel M. Génard, Bénédicte Quilot-Turion

► To cite this version:

Elsa Desnoues, Yves Y. Gibon, Valentina Baldazzi, Veronique Signoret, Michel M. Génard, et al.. High or low fructose? Consequences for sugar metabolism in peach fruit. 6. Rosaceous Genomics Conference, Sep 2012, San Michele all'Adige, Italy. 1 p., 2012. hal-02804340

HAL Id: hal-02804340

<https://hal.inrae.fr/hal-02804340>

Submitted on 5 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

High or low fructose? Consequences for sugar metabolism in peach fruit

E. Desnoves¹, Y. Gibon², V. Baldazzi³, V. Signoret¹, M. Génard³, B. Quilot-Turion¹

¹ INRA, UR1052 Génétique et Amélioration des Fruits et Légumes F-84143 Montfavet, France, ² INRA, UMR1332 Fruit Biology and Pathology F-33883 Villenave d'Ornon, France, ³ INRA, UR1115 Plantes et Systèmes de culture Horticoles F-84914 Avignon, France


Introduction

Fruit taste is largely affected by the content of sugars and acids. Fructose is the sweetest tasting sugar. In commercial peach fruit, sucrose is the main sugar, followed by fructose and glucose which have similar levels. Interestingly low fructose accessions have been described in wild peaches^{1,2}. Through an extensive profiling of metabolites and enzymatic activities, this study aims at i) describing sugar metabolism in peach fruit at different developmental stages and ii) comparing two genotypes with contrasted fructose/glucose ratios.


Materials and methods

Fruit samples were collected at 6 or 7 developmental stages respectively in 2010 and 2011. Three pools of 3 fruits were collected for each stage. Fruit mesocarp was cut in little pieces, immediately frozen in liquid nitrogen, ground to a fine powder and stored at -80°C. Twelve metabolites and 12 enzyme activities were measured by enzymatic methods³.

Hypotheses to explain 'low fructose' genotype : reduced synthesis and/or increased degradation



Results



Sugar metabolism in peach and time courses smoothed from concentration data of the different metabolites (mg/gFW), proteins and amino acids (mol/FW) along fruit growth (degree days after bloom) for 2 genotypes and 2 years.

SO	SDH	SuSy	SPS	F1,6bisP	NI	PFK	FRK	PGM	GK	AI	UGPase
0 - 0.1	0.2 - 0.7	1 - 3	1 - 5	10 - 60	20 - 80	10 - 150	50 - 150	60 - 150	60 - 200	50 - 800	450 - 600
→	→	→	→	→	→	→	→	→	→	→	→

Min and max activities (nmol/gFW/min) and mean tendency of evolution along fruit growth of 12 enzymes for 2 genotypes and 2 years

UGPase : UDP-glucose pyrophosphorylase; PGI : phosphoglucosyltransferase; PGM : phosphoglucosylmutase; PFP : phosphotransférase; PFK : phosphofructokinase; F1,6BisPase : fructose 1,6-bisphosphatase; SPS : sucrose phosphate synthase; FRK : fructokinase; GK : glucokinase; SuSy : sucrose synthase; NI : neutral invertase; AI : acid invertase; SDH : sorbitol dehydrogenase; SO : sorbitol oxidase

Sugar metabolism is a highly regulated system in which a major perturbation in a central compound has only slight repercussions on other metabolites and on enzyme activities

Other explanations for the low fructose phenotype

- **Differential affinities for substrate between iso-enzymes.** In peach fructokinase has two isoforms with different affinities for fructose².
- **Limited fructose storage resulting in higher degradation.** Fructose and glucose are stored in the vacuole. If fructose cannot be stored or if it 'destocked' to cytosol, it is more degraded than glucose.
- **Differential consumption of the two hexoses for respiration, cell wall or synthesis of other carbon compounds.** First results suggest that the 'low fructose' genotype has more carbon than the 'commercial' genotype though there is no difference between the two genotypes for dry mass.