



**HAL**  
open science

## Digestive anatomy and impact of dietary fibre on performances of the growing grasscutter (*Thryonomys swinderianus*)

Yapo Magloire Yapi, Francis Enjalbert, Thierry Gidenne, Daniel Zongo

### ► To cite this version:

Yapo Magloire Yapi, Francis Enjalbert, Thierry Gidenne, Daniel Zongo. Digestive anatomy and impact of dietary fibre on performances of the growing grasscutter (*Thryonomys swinderianus*). 15. International Congress of the European Society of Veterinary and Comparative Nutrition, Sep 2011, Saragosse, Spain. , 2011. hal-02804816

**HAL Id: hal-02804816**

**<https://hal.inrae.fr/hal-02804816>**

Submitted on 5 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# Digestive anatomy and impact of dietary fibre on performances of the growing grasscutter (*Thryonomys swinderianus*)

Y.M. Yapi<sup>1</sup>, F. Enjalbert<sup>2</sup>, T. Gidenne<sup>2</sup>, D. Zongo<sup>1</sup>

<sup>1</sup>National Polytechnic Institute Houphouët-Boigny of Yamoussoukro, Ivory Coast;

<sup>2</sup>INRA, INPT-ENSAT, ENVY, UMR 1289 Tandem, France

[f.enjalbert@envt.fr](mailto:f.enjalbert@envt.fr)

## Introduction

The grasscutter is a rodent herbivore recently domesticated in several countries of sub-Saharan Africa for meat production. However, the development of this production to a rational and more productive model needs a better knowledge of the digestive physiology of this animal. This work aimed to study the digestive anatomy of the growing grasscutter and determine the impact of dietary fibre on growth rate and health.



Photo 1. Growing Grasscutter

## Material and methods

In a first trial, 80 grasscutters (37 ± 5 days and 330 ± 70 g) were allocated to 4 groups of 20 animals and submitted to 4 diets containing 8, 14, 22 and 26% acid detergent fiber (ADF) on dry matter (DM) basis (ADF8, ADF14, ADF22 and ADF26). Individual weight and mortality were recorded for three months. In a second trial, 60 grasscutters (40 ± 7 days and 446 ± 152 g) were allocated to 3 groups of 20 animals, submitted to 3 diets containing 12, 17 and 22% ADF (ADF12, ADF17 and ADF22). Weight was measured for two months and pH, DM content and fresh weight of contents of digestive compartments were measured on animals slaughtered at 40, 83 and 103 days of age.

## Results and Discussion

In the first trial, 33% mortality was observed in ADF8 after 7 weeks due to hemorrhagic enteritis. No mortality was observed in the other groups. Growth was most rapid with diet ADF14 followed by diet ADF22 and diet ADF26 ( $P < 0.05$ ) (figure 1). In the second trial, growth was most rapid with diet ADF22 followed by diet ADF17 (Figure 2). DM content, pH and relative importance of the compartments (Table 1) did not vary with age or diet ( $P > 0.05$ ).




Figure 1. Evolution of animal weight in first trial




Figure 2. Evolution of animal weight in 2nd trial

**Table 1.** Some parameters of the digestive tract

Digestive compartments	Stomach	Small intestine	Caecum	Proximal colon	Distal colon
pH	2.1 ± 0.7	6.5 ± 0.6	6.0 ± 0.3	6.4 ± 0.3	-
DM content of digestive compartments (%)	27.8 ± 7.3	19.0 ± 5.5	20.7 ± 8.9	23.2 ± 11.8	33.1 ± 14.1
% of total fresh digestive content	23.6 ± 1.7	16.8 ± 1.3	44.0 ± 9.0	10.5 ± 0.8	5.1 ± 0.2

## Conclusion

The optimum dietary level of lignocellulose for the growing grasscutter would be between 14 and 22%. The caecum appears to play an important role in the digestive function of this animal with more than 40% of total contents of the digestive tract.