



Spatial organization of tobacco cell plasma membrane : characterization, and modulation upon elicitation

Patricia Gerbeau Pissot, Christophe Der, Iulia-Andra Anca, Kévin Grosjean, Dominique Thomas, Yann Roche, Jean-Marie Perrier-Cornet, Sébastien Mongrand, Françoise Simon Plas

► To cite this version:

Patricia Gerbeau Pissot, Christophe Der, Iulia-Andra Anca, Kévin Grosjean, Dominique Thomas, et al.. Spatial organization of tobacco cell plasma membrane: characterization, and modulation upon elicitation. Ecole Regulation et Dynamique des Membranes Biologiques, May 2013, Porquerolles, France. hal-02805522

HAL Id: hal-02805522

<https://hal.inrae.fr/hal-02805522>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Spatial organization of tobacco cell plasma membrane: characterization, and modulation upon elicitation

Patricia Gerbeau-Pissot¹, Christophe Der¹, Iulia Anca², Kevin Grosjean¹, Dominique Thomas¹, Yann Roche², Jean-Marie Perrier-Cornet³, Sébastien Mongrand⁴, Charles Kervrann⁵ and Françoise Simon-Plas²

1- INRA, Agroecologie, CNRS, France

2- Université de Bourgogne, CNRS, France

3- AgroSup Dijon, Laboratoire Procédés Alimentaires et Microbiologiques, France

4- CNRS, Laboratoire de Biogénèse Membranaire, Université Victor Segalen, INRA, France

5- INRIA Rennes- Bretagne Atlantique, Campus Universitaire de Beaulieu, France

The ability of lipids to form liquid-ordered sterol-rich phases in complex mixtures, and the associability of specific proteins with sterol-enriched biochemical fractions extracted from biological membranes, have given rise to the "lipid raft" hypothesis. This hypothesis posits the presence of small-sized ordered domains of particular lipid and protein composition within biological membranes, which could act as signal transduction platforms, particularly during plant- microorganism interactions. However, the precise organization of the living cell plasma membrane remains poorly described. For example, the spatial distribution of segregating liquid-disordered and liquid-ordered phases has not been analyzed in plant cells at a resolution compatible with the estimated size of such assemblies.

We developed a multispectral confocal microscopy approach to generate ratiometric images of a large surface of tobacco cell plasma membrane, suggesting the membrane exhibits a mosaic organization into areas of various levels of order. Moreover, we demonstrate that modification of the plasma membrane fluidity, together with an increase in the proportion and organization of ordered domains, transiently occurred in the early steps of signaling triggered by cryptogein, an elicitor of defense reaction.