



HAL
open science

Viability analysis of the natural population of *Salmo salar* L. in the Allier catchment : impact of 35 years stocking

Guillaume Dauphin, Etienne Prévost

► **To cite this version:**

Guillaume Dauphin, Etienne Prévost. Viability analysis of the natural population of *Salmo salar* L. in the Allier catchment : impact of 35 years stocking. Canadian Conference for Fisheries Research, Société Canadienne de Limnologie. CAN., Jan 2013, Windsor, Canada. 1 p. hal-02805529

HAL Id: hal-02805529

<https://hal.inrae.fr/hal-02805529>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Invited Oral SCL (Experimental Lakes Area Research)

VIABILITY ANALYSIS OF THE NATURAL POPULATION OF SALMO SALAR L. IN THE ALLIER CATCHMENT: IMPACT OF 35 YEARS OF STOCKING

Dauphin*^{1,2}, G. and E. Prévost^{2,3}

¹INRA, UMR 1224 ECOBIOP, Pôle d'hydrobiologie, Ibaron, 64310 Saint-Pée sur Nivelle, France. ²Science Branch, Department of Fisheries and Oceans, Northwest Atlantic Fisheries Center, 80 East White Hills Road, PO Box 5667 St John's NL, CANADA A1C 5X1 (Guillaume.Dauphin@dfo-mpo.gc.ca). ³UPPA, UMR ECOBIOP, UFR Sciences et Techniques Côte Basque, Campus Montauray, 64600 Anglet, France.

Due to its conservation value, the Allier's salmon population has been monitored and several datasets such as fisheries catches, spawning nest counts, juvenile index of abundance and the number and life-stage of salmon stocked every year have been collected from 1975 to present. Synthesizing the information brought by these heterogeneous data sources in a formal statistical modelling framework is a difficult task. In order to reflect the natural process governing the population renewal it is also important to account for regulation mechanism such as density dependence as well as variability in the different transition parameters such as survival. Hierarchical Bayesian modelling (HBM) offers an efficient way to deal with such constraints while accounting for various forms of uncertainty. The model built during this study and presented here brings together 35 years of heterogeneous data in a coherent framework while accounting for uncertainty. The results show a retrospective estimation of the past abundance of Atlantic salmon in three different spatial areas of the Allier River as well as the intergenerational renewal rate of the population. One of the main challenges of this modelling exercise was to incorporate the annual stocking data. The model provides estimates of the contribution of the different categories of salmon life-stage stocked (egg, fry and smolt) over the time series considered. These results provide useful information to the managers to apprehend the impact of the different restoration program over the last decades in the Allier River and make decision about the future programs.

Oral CCFR (Disturbed Ecosystems, Threatened Species and Restoration)

RECOVERY STRATEGY FOR A STRADDLING STOCK: 3LNO AMERICAN PLAICE

Dauphin*, G. and P. Shelton

Science Branch, Department of Fisheries and Oceans, Northwest Atlantic Fisheries Center, 80 East White Hills Road, PO Box 5667 St John's NL, CANADA A1C 5X1 (Guillaume.Dauphin@dfo-mpo.gc.ca).