



Carbon dynamics of intensively managed forest along a full rotation

Virginie Moreaux, Alexandre Bosc, Jean-Marc J.-M. Bonnefond, Régis R. Burlett, Eric Lamaud, Michel Sartore, Pierre Trichet, Christophe Chipeaux, Catherine Lambrot, Andrew S. Kowalski, et al.

► To cite this version:

Virginie Moreaux, Alexandre Bosc, Jean-Marc J.-M. Bonnefond, Régis R. Burlett, Eric Lamaud, et al.. Carbon dynamics of intensively managed forest along a full rotation. AGU Fall Meeting, Dec 2012, San Francisco, United States. n.p., 2012. hal-02805684

HAL Id: hal-02805684

<https://hal.inrae.fr/hal-02805684>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

carbon dynamics of intensively managed forest along a full rotation: a juvenile versus adult stage approach

MOREAUX V.¹, A. BOSCH¹, J.-M. BONNEFOND¹, R. BURLETT², E. LAMAUD¹, M. SARTORE¹, P. TRICHET¹, C. CHIPEAUX¹, C. LAMBROT¹, A. KOWALSKI³ & D. LOUSTAU¹
¹ UR 1263 EPHYSE - INRA, Villenave d'Ornon, France, ² UMR BIOGECO - INRA Talence, France, ³ Centro Andaluz de Medio Ambiente, Granada, Spain

Contact : virginie.moreaux@bordeaux.inra.fr

1. Introduction

- Almost all European forests are transformed by management [1] with varying intensities and silvicultural operations have profoundly modified the natural disturbance regime. Forest management exerts a major influence on the contribution of temperate forests to global carbon sequestration [2]. Meanwhile, it is well recognized that northern forests contain large amounts of carbon (C) in both biomass and soils and these reservoirs may be more vulnerable to future climate changes [1].
- Managed forests present mosaic of stand ages that are continuously being renewed through natural (drought, storm...) and anthropogenic disturbances (soil preparation, thinning, understorey control, harvest, shorter rotation...), which influence forest growth and the concomitant energy, water and carbon exchanges [3,4,5,6].
- They result ultimately in either positive or negative changes in the biomass and soil carbon pools but few, if any, datasets or modeling tools are available for quantifying these impacts.
- Accurate accounting of these large C stocks requires a sound scientific understanding of how disturbances and climate variability impact the emission and sequestration of carbon and how their respective effects might be factored out.

Objective: For disentangling these impacts, we have collected a long term series of half-hourly latent, sensible heat and CO₂ fluxes from a managed pine forest and we analysed the effect of management operations, tree age and climate using these data.

2. Experiments

Bilos (juvenile stage)

Le Bray (adult stage)

Eddy flux measurements,
Biomass measurements,
Soil measurements,
Meteorological measurements

3. Silvicultural operations




Figure 2. Evolution of the carbon fluxes for a characteristic meteorological year, in the young stand (blue lines) and in the adult one (red lines)

- Plowing and sowing operations increase the heterotrophic respiration through massive inputs of dead plant material: dead wood, organic humus layers and ground vegetation and the activation of the mineralization of soil organic matter.
- The succession of silvicultural operations during the site preparation made the ecosystem a net carbon source: 10 years after clearcutting, the cumulative balance is still positive, but 2 years after the sowing, the site rapidly becomes a sink of CO₂.
- The impact of thinning and vegetation removal (weeding) at the age of 5 years old brought the balance to almost neutral (2009).
- In the adult stage, no significant impact of silvicultural operation: strong carbon sequestration sensitive to climate

4. Age effects




Figure 3. Evolution of the carbon fluxes for a characteristic meteorological year, in the young stand (blue lines) and in the adult one (red lines)





Figure 1. Cumulative net ecosystem exchange calculated over an 8-year period in the young stand (blue lines) and in the adult one (red lines)

- Under characteristic meteorological conditions, the magnitude of the carbon fluxes increased with stand age. Annual GPP reached 2087 gC m⁻² against 881 gC m⁻² for the young stand, and the onset of the growing season is earlier in the adult stand. These observations are explained by the difference in soil available water, stand LAI being similar for the two sites (3.5 m² m⁻²).
- We noticed a higher exposure to soil drought in the young stand due to incomplete soil exploration by roots and thus lower soil water extractable.
- Understorey from the young stand is more sensitive to soil drought [6].

5. Climate effects



Our data show that the management operations masked the impact of soil water deficit at the juvenile stage.

- In both stands, a drastic decrease in GPP occurred during the summer dry year 2005 compared to the characteristic year 2008.
- In the adult stand, the drop in GPP occurred as SWC reached the wilting point (Bray, Figure 4 orange lines).
- In the young stand, carbon fluxes in 2005 are mostly explained by the effect of the previous ploughing: ecosystem respiration dominates and GPP remained low most part of the year.
- Therefore, the site is a carbon source all along the 2005 year.

Figure 4. Evolution of the carbon fluxes for a characteristic meteorological year (Green lines) and a dry year (Orange lines), on both sites.

6. Carbon stocks

Bilos (juvenile)			
Carbon stocks (gC m ⁻²)	2000	2010	C stock variations
Aboveground biomass	1455 ± 92	2170 ± 66	+ 715
Belowground biomass	1843 ± 15	528 ± 21	- 1315
Organic soil	2875 ± 206	291 ± 75	- 2585
Mineral soil	9881 ± 1093	12187 ± 1010	+ 2305
Total (gC m ⁻²)	16056 ± 1123	15176 ± 1171	- 880

Carbon balance is consistent between stocks and fluxes

Bray (mature)			
Carbon stocks (gC m ⁻²)	2003 [7]	2004 [8]	2009 [9]
Organic soil	1410 ± 304	3063 ± 175	
Mineral soil	10890 ± 1914	9917 ± 802	
Total (gC m ⁻²)	12300 ± 2218	12980 ± 977	

No significant changes

References

- [1] FAO 2010
 [2] Ciais et al. 2008
 [3] Granier et al. 2007
 [4] Sohngen 2008
 [5] Kowalski et al. 2004
 [6] Moreaux et al. 2011
 [7] Guillet et al. 2010
 [8] Schrupp et al. 2011
 [9] Schrupp, pers. Comm
 [10] Loustau et al. 2012.
 AGU poster n°B51B0492

Acknowledgements

This study was supported by the GHG-Europe project.
 Flux data collected at the young and adult pine sites were also obtained from D. Garrigou, M. Irvine, J. Ogée, S. Delzon.