



**HAL**  
open science

## **Decomposing sensible fluxes from two urban zones using LAS scintillometry over the city of Nantes during the FLUXSAP 2012 measurement campaign**

Mark Rankin M. R. Irvine, Jean-Pierre Lagouarde, Sylvia Dayau, Aurore Brut, Jean-Martial Cohard, Georges Najjar, Fabrice Rodriguez, Jean-Marc Rouaud, Patrice Mestayer

### **► To cite this version:**

Mark Rankin M. R. Irvine, Jean-Pierre Lagouarde, Sylvia Dayau, Aurore Brut, Jean-Martial Cohard, et al.. Decomposing sensible fluxes from two urban zones using LAS scintillometry over the city of Nantes during the FLUXSAP 2012 measurement campaign. Tübingen Atmospheric Physics Symposium "Scintillometers and Applications", Oct 2013, Tübingen, Germany. hal-02806369v1

**HAL Id: hal-02806369**

**<https://hal.inrae.fr/hal-02806369v1>**

Submitted on 6 Jun 2020 (v1), last revised 20 Jul 2020 (v2)

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## DECOMPOSING SENSIBLE FLUXES FROM TWO URBAN ZONES USING LAS SCINTILLOMETERY OVER THE CITY OF NANTES DURING THE FLUXSAP 2012 MEASUREMENT CAMPAIGN.

Irvine M.R. (1), Lagouarde J-P. (1), Dayau S. (1), Brut A. (2), J-L., Cohard J-M. (3), Najjar G. (4), Rodriguez F. (5), Rouaud J-M. (5) Mestayer P. (6)

- (1) INRA, UR1263 EPHYSE, F-33140 Villenave d'Ornon
- (2) CESBIO, UMR5126, F-31401 Toulouse
- (3) LTHE, UMR5564, F-38400 Saint-Martin d'Hère
- (4) LSIIT, UMR7005, F-67412 Illkrich
- (5) IFSTTAR, GER, F-44341
- (6) IRSTV, FR CNRS 2488, F-44321

In the framework of a recent research program dealing with the sustainable development of cities (ANR 'Villes Durables', 'VegDUD'), two intensive field experiments (FluxSAP, <http://www.irstv.fr/> Mestayer [1]) were installed over the city of Nantes, in France. After an initial experiment in 2010, a second experiment in 2012 was set up to investigate different urban zones within Nantes. One an artificial surface (a parking lot), a typical 1960's urban housing zone and a modern 'eco-quarter' housing zone built in 2010/11. The experiment ran during the month of June 2012. As part of the set-up 4 Large Aperture Scintillometers (LAS) were installed over the city with distances ranging from 800m to 3km. The installation was set-up in such a way that there is one trajectory, which integrates roughly over all surfaces, with the others installed to decompose the long trajectory into smaller 'typical' surfaces.

Sensible heat flux along each path will be presented in conjunction with footprint estimates developed using the method from Meijninger [2] based on the dispersion model of Horst [3]. Interpretation of the results with regard to surface type will be discussed.

### References

- [1] Mestayer, P.G., Rosant, J.-M., Rodriguez, F., and Rouaud, J.-M., 2011, La Campagne Expérimentale FluxSAP 2010, La Météorologie 73, pp. 34-44
- [2] Meijninger, W.M.L., Hartogensis, O.K., Hoedjes, J.C.B., Zuurbier, R.M. and DeBruin, H.A.R. 2002, Determination of Area-Averaged Sensible Heat Fluxes with a Large Aperture Scintillometer over a Heterogeneous Surface, Bound. Layer Meteorol. 105, pp. 37-62
- [3] Horst, T.W. and Weil, J.C., 1992, Footprint Estimation for Scalar Flux Measurements in the atmospheric surface-layer, Bound. Layer Meteorol. 59, pp. 279-296.