



HAL
open science

A multi-instruments campaign using scintillometry to measure the horizontal turbulence spatial distribution

Clélia Robert, Jean-Martial Cohard, Jean-Pierre Lagouarde, Mark Rankin M. R. Irvine, Aurore Brut

► **To cite this version:**

Clélia Robert, Jean-Martial Cohard, Jean-Pierre Lagouarde, Mark Rankin M. R. Irvine, Aurore Brut. A multi-instruments campaign using scintillometry to measure the horizontal turbulence spatial distribution. Tübingen Atmospheric Physics Symposium "Scintillometers and Applications", Oct 2013, Tübingen, Germany. hal-02806387

HAL Id: hal-02806387

<https://hal.inrae.fr/hal-02806387>

Submitted on 6 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

A multi-instruments campaign using scintillometry to measure the horizontal turbulence spatial distribution

C. Robert (1), J. M. Cohard (2), J.P. Lagouarde (3), M. Irvine (3) and A. Brut (4).

(1) ONERA, The French Aerospace Lab France.

(2) LTHE, UMR5564, Univ. de Grenoble, France

(3) INRA, UR1263 EPHYSE, 71 avenue E. Bourlaux, F-33140 Villenave d'Ornon

(4) CESBIO, UMR 5126 Université Paul Sabatier, 18 Avenue E. Belin 31401 Toulouse Cedex 9

Agregated fluxes measurements with scintillometric methods rely on horizontal homogeneity assumptions (Monin-Obukov theory). If these measurements were shown valid over slightly heterogeneous terrain [1], questions still remain when spatial heterogeneity is more pronounced (leading to Bowen ratio variability) which is susceptible to generate steady turbulent structures with a large range of C_n^2 values [2].

A field experiment has been conducted near Lannemezan (France) aiming to evaluate new scintillometric methods to quantify the spatial variability of refractive index structure parameter (C_n^2) from which heat fluxes can be derived. The C_n^2 variability comes from the intrinsic topography along the path. Three devices were installed between the Centre de Recherche Atmosphérique and the Campistrous church, 2.7 km northward: The Shack-Hartmann wavefront sensor (SH) working in 3.8-4.2 μm wavelengths [3]. It consists in saving a matrix of 25 sub-pictures of a double source from which we compute cross correlations of the scintillation indices and angles of arrival. 5 to 8 C_n^2 values along the path can be extracted from these correlations. 5 scintillometers with different aperture sets measured an averaged C_n^2 for different parts along the path. Moreover 2 eddy-covariance stations (EC) measured the local structure parameters C_T^2 , C_Q^2 et C_{TQ} to calculate the local C_n^2 . Continuous time series of C_n^2 measurements from the scintillometers have been stored together with EC measurements from 28/08/2012 till 22/10/2012. From the 14/09/2012, 21 days and 5 nights data sequences have been stored with the SH. Data analysis of the SH measurements produced spatial and temporal spectra of scintillations and of angles of arrival that show a good concordance to the Kolmogorov model. We checked that both sources are embedded in the same small turbulence regime. Then inversion of the SH measurements covariances should allow us to retrieve the $C_n^2(x)$ profile along the sightline. These local estimations will be compared with the C_n^2 integrated from the scintillometers.

References

[1] Lagouarde, J. P., Bonnefond, J. M., Kerr, Y. H., McAneney, K. J., and Irvine, M. (2002). Integrated sensible heat flux measurements of a two-surface composite landscape using scintillometry. *Boundary Layer Meteorology* 105, 5-35.

[2] Guyot, A., J.-M. Cohard, S. Anquetin and S. Galle (2012). "Long-term observations of turbulent fluxes over heterogeneous vegetation using scintillometry and additional observations: A contribution to AMMA under Sudano-Sahelian climate." *Agricultural and Forest Meteorology* **154–155**(0): 84-98.

[3] Robert, C., V. Michau, B. Fleury, S. Magli and L. Vial (2012). "Mid-infrared Shack-Hartmann wavefront sensor fully cryogenic using extended source for endoatmospheric applications." *Opt. Express* **20**(14): 15636-15653