



**HAL**  
open science

## **The ruminal level of trans-10 fatty acids of dairy cows is linked to the composition of bacterial community**

Asma Zened, Sylvie Combes, Laurent Cauquil, Jérôme J. Mariette, Chloé Rousseau, Christophe Klopp, Annabelle Troegeler-Meynadier, Francis Enjalbert

### ► **To cite this version:**

Asma Zened, Sylvie Combes, Laurent Cauquil, Jérôme J. Mariette, Chloé Rousseau, et al.. The ruminal level of trans-10 fatty acids of dairy cows is linked to the composition of bacterial community. 4. Congress of European Microbiologists, FEMS, Jun 2011, Genève, Switzerland. 1p, 2011. <hal-02806622>

**HAL Id: hal-02806622**

**<https://hal.inrae.fr/hal-02806622v1>**

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

# The ruminal level of *trans*-10 fatty acids of dairy cows is linked to the composition of bacterial community

A. ZENED<sup>1</sup>, S. COMBES<sup>1</sup>, L. CAUQUIL<sup>1</sup>, J. MARIETTE<sup>2</sup>, C. ROUSSEAU<sup>3</sup>, C. KLOPP<sup>2</sup>, A. TROEGELER-MEYNADIER<sup>1</sup>, F. ENJALBERT<sup>1</sup>  
<sup>1</sup>UMR 1289 Tandem, INRA / INPT-ENSAT / ENVT, Castanet-Tolosan, France, France <sup>2</sup>Plateforme bio-informatique Genotoul, Biométrie et Intelligence Artificielle, INRA, Castanet-Tolosan, France, <sup>3</sup>INPT-ENSAT, Castanet-Tolosan, France  
 a.zened@envt.fr

## Objective and Methods




Fig.1 Biohydrogenation of linoleic acid (C18:2)

In dairy cows, *trans*-10 fatty acids (*t10*-FA) are produced by ruminal bacteria during lipid digestion, in particular during C18:2 biohydrogenation (Fig. 1), thereafter absorbed and secreted into milk (Fig. 2).

High *t10* FA production causes milk fat depression and is detrimental for human consumer health.





Fig. 2

This study investigated the relationship between the ruminal proportion of *t10* FA and the composition of rumen bacterial community in a dairy cow herd.


Ruminal fluids of 10 cows receiving the same diet and exhibiting a high heterogeneity of *t10*-FA ruminal proportions

DNA extraction




Barcoded reads of the V3-V4 regions of the 16S rRNA genes, were obtained from a 454 GS-FLX sequencer

Taxonomic assignments were performed using the open-source software Mothur V.1.12.3 ([www.mothur.org](http://www.mothur.org))



## Results and conclusions

A negative correlation ( $R=-0.7$ ) was noticed between *t10* FA and milk fat content (MFC) which was, on the contrary, positively but poorly correlated ( $R=0.2$ ) to *t11* FA.



Variation in ruminal percentage of *t10* FA was associated with changes in the ruminal bacterial community

A relationship between lactic acid production and high levels of *t10* FA has already been observed. In this study, an increase in Veillonellaceae family and *Lactobacillus* genus, which are consumers and producers of lactic acid respectively, is noticed in ruminal fluids exhibiting high *t10* FA percentage.

The abundance of other bacteria were also linked to *t10* FA production but their functions are not yet well-established.

Phylum	Order	Family	Genus	R( $\geq 0.7$ )	P-value	Abundance (%)
Firmicutes	Clostridiales	Lachnospiraceae		0.8	***	18.05
Firmicutes	Clostridiales	Lachnospiraceae	<i>Syntrophococcus</i>	0.9	***	0.87
Firmicutes	Clostridiales	Lachnospiraceae	<i>Butyrivibrio-Pseudobutyrvibrio</i>	-0.8	**	0.18
Firmicutes	Clostridiales	Veillonellaceae	<i>Dialister</i>	0.7	**	0.97
Firmicutes	Lactobacillales	Lactobacillaceae	<i>Lactobacillus</i>	0.8	***	0.14
Bacteroidetes	Bacteroidales	Prevotellaceae	uncultured	-0.7	*	1.91
Bacteroidetes	Bacteroidales	RF16		-0.7	**	0.70
Actinobacteria	Coriobacteriales	Coriobacteriaceae		0.9	***	0.80

\* <0.05  
 \*\* <0.01  
 \*\*\* <0.001