



**HAL**  
open science

## **Attenuation of brown adipose tissue activity during chemotherapy treatment: a new mechanism of weight gain in breast cancer women?**

Emilie E. Gadéa, E. Thivat, C. Merlin, R. Paulon, F. Kwiatkowski, J.B Chadeyras, B. Coudert, Yves Y. Boirie, Béatrice Morio, X. Durando

### ► To cite this version:

Emilie E. Gadéa, E. Thivat, C. Merlin, R. Paulon, F. Kwiatkowski, et al.. Attenuation of brown adipose tissue activity during chemotherapy treatment: a new mechanism of weight gain in breast cancer women?. Obesity, Physical Activity and Cancer Conference 2013, World Cancer Research Fund International (WCRF). Londres, GBR., Apr 2013, Londres, United Kingdom. hal-02806643

**HAL Id: hal-02806643**

**<https://hal.inrae.fr/hal-02806643v1>**

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Attenuation of brown adipose tissue activity during chemotherapy treatment: a new mechanism of weight gain in breast cancer women?

Gadea E.<sup>1,2,3</sup>, Thivat E.<sup>1,4</sup>, Merlin C.<sup>1</sup>, Paulon R.<sup>1</sup>, Kwiatkowski F.<sup>1</sup>, Chadeyras J-B.<sup>1</sup>, Coudert B.<sup>5</sup>, Boirie Y.<sup>2,6</sup>, Morio B.<sup>2</sup> and Durando X.<sup>1,3,7</sup>

<sup>1</sup>Centre Jean Perrin, Clermont-Ferrand, France <sup>2</sup>Clermont University, University of Auvergne, UNH UMR1019, Clermont-Ferrand, France <sup>3</sup>CIC, Clermont-Ferrand, France <sup>4</sup>EA 4677 ERTICA, Clermont-Ferrand, France <sup>5</sup>Centre Georges-François Leclerc, Dijon, France <sup>6</sup>CHU Clermont-Ferrand, Clermont-Ferrand, France <sup>7</sup>Translational oncology EA3846 University of Auvergne, Clermont-Ferrand, France

**Background and Aims:** Weight gain has been reported in 17 to 34% of early stage breast cancer patients during chemotherapy. This gain has become a major concern because it has been suggested that weight variation was associated to a poor prognosis. However the mechanisms involved remain unclear. Alteration in thermogenesis induced by a decrease of brown adipose tissue (BAT) activity after chemotherapy, may partly contribute to weight gain in these patients.

**Methods:** A PET/CT scan was performed at baseline and after one course of docetaxel + trastuzumab treatment in 26 breast cancer women. Variation of 18F-FDG uptake in BAT between the two measures was assessed retrospectively according to weight changes.

**Results:** Percentage of patients who gained weight was 35%, 25% lost weight and 40% remained stable. Women who gained weight during chemotherapy experienced a significant decrease of 18F-FDG uptake in BAT ( $p=0.0048$ ).

**Conclusion:** These original data suggest for the first time that BAT modulation by chemotherapy would be a potential contributor to body weight gain through blunted thermogenesis in breast cancer patients.