



## The identification of a new genotype of avian paramyxoviruses type I in West-Africa provides new outcomes for phylogeny reconstruction

Saliha Hammoumi, Renata Servan de Almeida, Patricia Gil, F. Briand, Sophie Molia, Nicolas Gaidet, J. Cappelle, Véronique Chevalier, Gilles Balança, A. Traore, et al.

### ► To cite this version:

Saliha Hammoumi, Renata Servan de Almeida, Patricia Gil, F. Briand, Sophie Molia, et al.. The identification of a new genotype of avian paramyxoviruses type I in West-Africa provides new outcomes for phylogeny reconstruction. EPIZONE 5th Annual Meeting, Apr 2011, Arnhem, Netherlands. pp.1, 2011. hal-02806752

**HAL Id: hal-02806752**

**<https://hal.inrae.fr/hal-02806752>**

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# The identification of a new genotype of avian paramyxoviruses type I in West-Africa provides new outcomes for phylogeny reconstruction

S. Hammoumi<sup>1</sup>, R. Servan de Almeida<sup>1</sup>, P. Gil<sup>1</sup>, F. Briand<sup>2</sup>, S. Molia<sup>3</sup>, N. Gaidet<sup>3</sup>, J. Cappelle<sup>3</sup>, V. Chevalier<sup>3</sup>, G. Balança<sup>3</sup>, A. Traoré<sup>4</sup>, C. Grillet<sup>1</sup>, K. Samaké<sup>4</sup>, A. Diarra<sup>4</sup>, D. Martinez<sup>1</sup>, V. Jestin<sup>2</sup>, E. Albina<sup>1</sup>



<sup>1</sup> CIRAD, UMR CIRAD/INRA CMAEE, Montpellier, France.

<sup>2</sup> Anses-Ploufragan-Plouzané VIPAC, French Reference Laboratory for Avian Influenza and Newcastle Disease, Ploufragan, France

<sup>3</sup> CIRAD, UPR AGIRS, Montpellier, France.

<sup>4</sup> Laboratoire Central Vétérinaire, Bamako, Mali.




## INTRODUCTION




Newcastle disease (ND) is one of the most severe infectious diseases of birds, particularly poultry, and has been the cause of serious economic losses worldwide. Newcastle disease virus (NDV) or avian paramyxovirus type 1 (APMV-1) are divided in two clades (class I and class II) according to the genome size as well as sequence of the F and L genes. Virulent strains are found in the class II. There is no clear consensus for the NDV class II taxonomy since some authors use genotypes [1] whereas others use lineages, both covering distinct isolate clusters. Ten genotypes were traditionally considered for class II but a 11<sup>th</sup> genotype was recently proposed by our group. Aldous et al. [2] proposed 6 phylogenetic lineages based on the topology of the maximum likelihood tree of the 375bp of the 3' end of the F gene (further subdivided into 13 sublineages) to replace the genotype nomenclature previously adopted [1].

In many parts of Africa, ND is considered as endemic but only few data are available on the variants involved. In West Africa, new subgenotypes VII<sub>f</sub>, VII<sub>g</sub> and VII<sub>h</sub> were recently described [3] and integrated into the lineage 5 described by Aldous et al. [2]. Based on phylogenetic analyses of a partial F coding sequence of NDV isolates recovered from apparently healthy poultries in Mali in 2007 and 2008, we previously proposed a new subgenotype VII<sub>i</sub> [4]. Cattoli et al [5] have also described new strains that were clustered into a new lineage 7 and four new sublineages, 7a, 7b, 7c and 7d, corresponding to our subgenotypes VII<sub>i</sub>, VII<sub>g</sub>, VII<sub>h</sub> and VII<sub>f</sub>, respectively.




**Figure 1. schematic organisation of Newcastle disease virus genome and viral particle.**

N: nucleocapsid, P: phosphoprotein, M: matrix, F: fusion, HN: hemagglutinin-neuraminidase, L: RNA-directed RNA polymerase. Two additional proteins (V and W) are encoded by the RNA editing of the P gene.



In this study we present and describe new NDV detected in West-Africa. Sequences from these viruses and others publicly available were included in an extensive phylogeny reconstruction with different methods. We propose a refined taxonomy for NDV genotyping showing better consistency than the former lineage nomenclature. Particularly, we introduce two new genotypes, XII and XIII using the analyses of the full length amino acids sequence of F and HN proteins.


## MATERIALS AND METHODS



**Figure 2. complete procedure for new NDV strains analysis**

## RESULTS


Ancient nomenclature	New nomenclature
Lineage 7	Genotype XIII
Genotype VIIc, d, e	Genotype XII
Genotype VIIb	Genotype VII



**Figure 3. Phylogenetic analysis of 85 complete F and HN protein sequences of APMV-1.** Trees were constructed using the maximum likelihood method with 1000 resampling for the bootstrap analysis. A class 1 strain was used as an outgroup. The F and HN trees were identical, thus, only the F tree is represented, where informative, branch support values for HN are indicated after the F bootstrap values. Sequences generated by our group are indicated with a parallelogram.

## REFERENCES

- Ballagi-Pordany A, Wehmann E, Herczeg J, Belak S, Lomniczi B (1996). Identification and grouping of Newcastle disease virus strains by restriction site analysis of a region from the F gene. Arch Virol 141: 243-261.
- Aldous EW, Myn JK, Banks J, Alexander DJ (2003). A molecular epidemiological study of avian paramyxovirus type 1 (Newcastle disease virus) isolates by phylogenetic analysis of a partial nucleotide sequence of the fusion protein gene. Avian Pathol 32: 239-256.
- Snoeck CJ, Ducatez MF, Owoade AA, Faleke OO, Alkali BR, et al. (2009). Newcastle disease virus in West Africa: new virulent strains identified in non-commercial farms. Arch Virol 154: 47-54.
- Servan de Almeida R, Maminiaina OF, Gil P, Hammoumi S, Molia S, et al. (2009). Africa, a reservoir of new virulent strains of Newcastle disease virus? Vaccine 27: 3127-3129.
- Cattoli G, Fusaro A, Monne I, Molia S, Le Menach A, et al. (2009). Emergence of a new genetic lineage of Newcastle disease virus in West and Central Africa-Implications for diagnosis and control. Vet Microbiol 19: 142(3-4): 168-76.
- Abolnik C, Gerdes GH, Kitching J, Swanepoel S, Romito M, et al. (2008). Characterization of pigeon paramyxoviruses (Newcastle disease virus) isolated in South Africa from 2001 to 2006. Onderstepoort J Vet Res 75: 147-152.
- Herczeg J, Wehmann E, Bragg RR, Travassos Dias PM, Hadjiev G, et al. (1999). Two novel genetic groups (VIIb and VIII) responsible for recent Newcastle disease outbreaks in Southern Africa, one (VIIb) of which reached Southern Europe. Arch Virol 144: 2087-2099.



**Figure 4. Comparison of NDV genotyping using four methods of phylogeny reconstruction: maximum likelihood (ML), Bayesian (Mr Bayes), maximum parsimony (MP) and neighbour joining (NJ).** A total of 85 complete sequences of the F protein representing thirteen different genotypes were used in the phylogenetic reconstruction. Branch support values correspond to 1000 bootstrap replicates for ML, MP and NJ, and posterior probabilities estimated for 10,000 samples of the Markov chain for Mr Bayes. Area of the triangle is proportional to the number of isolates within the corresponding genotype.

## DISCUSSION AND CONCLUSION

In this study, a comprehensive phylogenetic reconstruction of NDV was carried out. The data sets used and the comparison of different methods for tree reconstruction on the F and HN proteins allow us to refine the NDV classification. We consider that the genotype nomenclature proposed by Ballagay-Pordany [1] is more appropriate to describe phylogenetic relationships between isolates, including the new isolates from West-Africa or Madagascar. Indeed, the clustering of the different genotypes supported by the analyses of the minimal distances between and within genotypes (not shown) do not match with the lineage nomenclature proposed by Aldous et al [2] and adopted by others [3, 5, 6].

Our conclusion is supported by the agreement in the phylogeny reconstruction obtained by four different methods on the complete F protein sequence. The resulting trees were all very similar but statistical analysis clearly shows that the ML and Bayesian methods are more robust.

From our work, the subgenotype VII<sub>b</sub> proposed by Herczeg [7] forms a distinct cluster in the branch including subgenotypes VII<sub>a</sub>, c, d, e. Therefore, we propose to limit the genotype VII to the subgenotype VII<sub>b</sub> which is considered as the first appeared in the evolutionary model and to group subgenotypes VII<sub>a</sub>, c, d, e in a new genotype XII.

Similarly, our five Malian strains form a new cluster containing other strains from West-Africa described by Snoeck et al [3] and Cattoli et al [5]. Snoeck et al [3] described three new subgenotypes (f, g, h) of lineage 5 (genotype VII) circulating in West Africa. Our Malian strains c\_ML007\_08 and c\_ML029\_07 clustered in two of these subgenotypes (VII<sub>g</sub> and VII<sub>h</sub>, respectively). However, the other Malian strains g\_ML038\_07 and c\_ML225\_08 are distant enough to constitute a new clade corresponding to the lineage 7a proposed by Cattoli et al. [5]. We propose to introduce the new genotype XIII for this group of West-African strains. Inside the genotype XIII, four subdivisions are clearly identified as subgenotypes XIII<sub>a</sub> (ex lineage 7a), XIII<sub>b</sub> (ex subgenotype VII<sub>g</sub>), XIII<sub>c</sub> (ex subgenotype VII<sub>h</sub>) and XIII<sub>d</sub> (ex subgenotype VII<sub>f</sub>).

This phylogenetic analysis suggests that this genotype XIII has evolved from a common ancestor with genotype VII (previously named subgenotype VII<sub>b</sub>) and XII (previously named VII<sub>c</sub>, d, e) but in a different direction than described earlier [3,5]. This suggests that an original strain older than genotype VII, probably linked to genotype VI was introduced in West Africa and since then has evolved in an independent manner to give genotypes VII and XIII whereas another lineage led to subgenotypes XII mainly found in Asia.



Département BIOS

Centre de coopération internationale en recherche agronomique pour le Développement

Contact person: renata.servan\_de\_almeida@cirad.fr